

Anatomy and relationships of the fuxianhuiid euarthropod *Guangweicaris* from the early Cambrian Guanshan Biota in Kunming, Yunnan, Southwest China revisited

Yichen Wu and Jianni Liu *Acta Palaeontologica Polonica* 64 (3), 2019: 543-548 doi:https://doi.org/10.4202/app.00542.2018

The Guanshan Biota (Cambrian Series 2, Stage 4) is a typical Burgess Shale-type biota, and is one of the most significant Cambrian Konservat-Lagerstätten from China, in addition to the well-known Chengjiang and Kaili biotas. *Guangweicaris* is a relatively common arthropod from the Guanshan Biota. Based on new specimens from the Longbaoshan and Xinglongcun sections of the Wulongqing Formation in Kunming, we herein report novel features of *Guangweicaris spinatus*, such as the eye-bearing anterior sclerite that is associated with the head shield, as well as revised details of the trunk, including the ventral appendages. A detailed comparison of *Guangweicaris*, *Fuxianhuia* and *Shankouia* suggests a close affinity between *Guangweicaris* and *Fuxianhuia*, which improves our understanding of the evolution of early Cambrian euarthropods.

Yichen Wu [wuyicheneli@126.com] and Jianni Liu [liujianni@126.com], State Key Laboratory of Continental Dynamics, Shaanxi Key Laboratory of Early Life and Environment, Department of Geology, Northwest University, Xi'an, China.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see <u>creativecommons.org</u>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.