

Growth dynamics and body size evolution of South American long-necked chelid turtles: A bone histology approach

Maria Eugenia Pereyra, Paula Bona, Ignacio Alejandro Cerda, Juan Marcos Jannello, Marcelo Saúl De La Fuente, and Bárbara Desántolo

Acta Palaeontologica Polonica 65 (3), 2020: 535-545 doi:<https://doi.org/10.4202/app.00702.2019>

Among turtles, cases of “gigantism” occur mostly in pleurodiran Pelomedusoides and cryptodirans, but are infrequent among pleurodiran chelids, which are mostly small-medium sized turtles. *Yaminuechelys* spp. are extinct South American long-necked chelids (from the Late Cretaceous–early Paleocene of Patagonia, Argentina) with caparaces almost three times larger than their extant sister taxon, *Hydromedusa tectifera*. Since evolutionary changes in size can be analyzed based on growth dynamics, we studied growth strategies from an osteohistological point of view. We sampled both extinct (*Yaminuechelys maior*) and extant (*H. tectifera*) species, in order to test hypotheses related to the mechanisms involved in the macroevolution of size within this clade. For this purpose, thin sections of long bone (humerus and femur) shafts of specimens of different ontogenetic stages for these species were prepared. The osteohistological study reveals a similar growth dynamic in both taxa, with a poorly vascularized cortex dominated by parallel- fibered bone and interrupted by lines of arrested growth (LAGs). The huge body size of *Y. maior* appears to be a consequence of the prolongation of the growth phase, suggesting that it had a longer lifespan than *H. tectifera*, allowing to reach greater sizes. In this way, and assuming that there is no displacement at the beginning of development (e.g., a delay in the earliest stages of growth) in *H. tectifera*, the acquisition of a large size in *Yaminuechelys* would be explained by hypomorphosis of the former or hypermorphosis of the latter, depending on the reconstruction of the ancestral condition of this clade.

Key words: Testudines, Chelidae, growth rate, body size, paleohistology, ontogeny, Paleocene, Argentina

Maria Eugenia Pereyra [m.eugenia.pereyra@gmail.com] and Paula Bona [paulabona26@gmail.com], División Paleontología Vertebrados, Museo de La Plata (Unidad de Investigación Anexo), Facultad de Ciencias Naturales y Museo, Paseo del Bosque s/n, CP1900, La Plata, Buenos Aires, Argentina. Ignacio Alejandro Cerda [nachocerda6@gmail.com], CONICET; Instituto de Investigaciones en Paleobiología y Geología, Universidad Nacional de Río Negro y Museo Carlos Ameghino, Belgrano 1700, Paraje Pichi Ruca (predio Marabunta),

CP8300, Cipolletti, Río Negro, Argentina. Juan Marcos Jannello [marcosjannello@hotmail.com] and Marcelo Saúl De la Fuente [mdelafuente1910@gmail.com], Instituto de Evolución, Ecología Histórica y Ambiente (IDEVEA-CONICET, Universidad Tecnológica Nacional, FRSR) Av. Gral. J.J. Urquiza 314, CP5600, San Rafael, Mendoza, Argentina. Bárbara Desántolo [bdesantolo@med.unlp.edu.ar], Cátedra de Citología, Histología y Embriología A, Facultad de Ciencias Médicas, Universidad Nacional de La Plata, 60 y 122 s/n, La Plata, Buenos Aires, Argentina.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(1,625.8 kB\)](#) |

 [Supplementary file \(647.4 kB\)](#)