

## Late Triassic spinicaudatan crustaceans from southwestern Poland

Ewa Olempska

*Acta Palaeontologica Polonica* 49 (3), 2004: 429-442

An assemblage of well-preserved spinicaudatan crustaceans ('conchostracans') is described from lacustrine late Carnian claystone at Krasiejów in southwestern Poland. Their shell microstructure is similar to that in extant spinicaudatans. Five species identified there are assigned to the genera *Laxitextella*, *Menucoestheria* (first record in the European Triassic), and *Menucoestheria bocki* sp. nov. and *Krasiestheria parvula* gen. et sp. nov. are erected. Specimens of *Laxitextella laxitexta* are the most abundant, comprising nearly half of the assemblage, those of *Menucoestheria bocki* one quarter, *Laxitextella* sp. A, *Menucoestheria* ? sp., and *Krasiestheria parvula* form a minor component. Their ecological setting was probably similar to Recent relatives: temporary ponds of fresh water. The Late Triassic fauna in the German part of the same basin is closely similar to that in Poland. In Europe, the stratigraphic range of *Laxitextella laxitexta* is limited to the Middle Keuper (middle-late Carnian).

**Key words:** Crustacea, Spinicaudata, taxonomy, palaeoecology, shell structure, Triassic, Poland.

Ewa Olempska [[olempska@twarda.pan.pl](mailto:olempska@twarda.pan.pl)], Instytut Paleobiologii PAN, ul. Twarda 51/55, PL-00-818 Warszawa, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



[Full text \(2,035.5 kB\)](#)