

New teeth of allotherian mammals from the English Bathonian, including the earliest multituberculates

Percy M. Butler and Jerry J. Hooker

Acta Palaeontologica Polonica 50 (2), 2005: 185-207


New material and additional morphological details of a rare and phylogenetically significant large fossil snake, *Wonambi naracoortensis* Smith, 1976, are described from Pleistocene and Pliocene cave deposits in southern South Australia. The new data refute some previous interpretations of the morphology of this species, and have implications for the phylogenetic position of *Wonambi* relative to extant snakes and other fossils, including other Madtsoiidae. The nature of contacts among palatal, braincase, snout, and mandibular elements imply similar functional attributes to those of extant anilioid snakes: maxillae from multiple individuals show corrugated contact surfaces for the prefrontal, implying a tight suture; structures on the anterior and medial surfaces of the palatine choanal process are interpreted as forming extensive contacts with the vomer and parasphenoid; and the distinctly bounded facets on the basiptyergoid processes and pterygoid imply little or no capacity for anteroposterior sliding of the palatopterygoid arch, hence absence of the macrostomatan 'ptyergoid walk'. On the frontal, interolfactory pillars were either absent or very slender, and a deep, sculptured contact surface for the nasal implies a prokinetic joint was also absent. Margins of the frontal and parietal indicate broad entry of the sphenoid into the ophthalmic fenestra, as in *Dinilysia*. Similarity of elements and features of the braincase (trigeminal foramen, ear region, and basiptyergoid processes) with both lizards and extant snakes show that differences between snakes and other squamates have sometimes been overstated. The case for macrostomatan affinities of *Wonambi* is not supported by new evidence.

Key words: Allotheria, "Haramiyida", Multituberculata, dentition, occlusion, tooth cusps, Jurassic.

Percy M. Butler [percy@butler92.freemove.co.uk], Department of Biology, Royal Holloway, University of London, Egham, Surrey, TW20 0EX, United Kingdom;

Jerry J. Hooker [j.hooker@nhm.ac.uk], Department of Palaeontology, The Natural History Museum, Cromwell Road, London, SW7 5BD, United Kingdom.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(1,139.7 kB\)](#)