

## A review of the endocerid cephalopod *Protocyptendoceras* from the Floian (Lower Ordovician) of the Eastern Cordillera, Argentina.

Marcela Cichowolski

*Acta Palaeontologica Polonica* 54 (1), 2009: 99-109 doi:<http://dx.doi.org/10.4202/app.2009.0110>

Ordovician cephalopods from Western Gondwana are still poorly known and published data is in need of systematic and stratigraphical revision. In this work, the endocerid proterocameroceratid *Protocyptendoceras* from the Floian of the Eastern Cordillera of Argentina is revised using new material from La Ciénaga, 5 km upstream from Purmamarca. The species *P. corvalani* and *P. teichert* are considered to be junior synonyms of *P. fuenzalidae*, which is redescribed in detail and reillustrated. Specimens preserve adapical parts of isolated siphuncles, related here to the presence of endocones in a posterior portion. Evidently *P. fuenzalidae* had a nekto-benthic mode of life in a shallow water environment, oriented horizontally. Its palaeobiogeographical affinities are mainly with Floian proterocameroceratids from Eastern Gondwana, such as *Anthoceras decorum* and the related genera *Lobendoceras*, *Ventroloboceras*, and *Notocycloceras*.

**Key words:** Cephalopoda, Endocerida, Proterocameroceratidae, Floian, Ordovician, Eastern Cordillera, Argentina.

Marcela Cichowolski [[mcicho@gl.fcen.uba.ar](mailto:mcicho@gl.fcen.uba.ar)], Área de Paleontología, Departamento de Ciencias Geológicas, Universidad de Buenos Aires, Ciudad Universitaria, Pab. II, Intendente Güiraldes 2160 C1428EGA, Buenos Aires, Argentina. CONICET.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

