

## Commensalism in the fossil record: Eunicid polychaete bioerosion on Pliocene solitary corals

Jordi Martinell and Rosa Domènech

*Acta Palaeontologica Polonica* 54 (1), 2009: 143-154 doi:<http://dx.doi.org/10.4202/app.2009.0115>

Some solitary caryophylliid (*Caryophyllia*, *Trochocyathus*, and *Ceratotrochus*) and flabellid (*Flabellum*) scleractinian corals from Pliocene of Western Mediterranean exhibit long groove-shaped bioerosional structures running along the surface of the thecae. They are epigenic structures produced by an episkeletozoan and therefore, they are described as Fixichnia. Here we propose *Sulcichnus* as a new ichnogenus, with three new ichnospecies (*Sulcichnus maeandriiformis*, *S. helicoidalis*, and *S. sigillum*) to name this traces. *Sulcichnus* is attributed to the activity of polychaetes. Similar structures are recently produced by *Lumbrineris flabellicola*, a symbiotic eunicid which maintains a commensalistic relation ship with solitary corals. In the fossil record, *Sulcichnus* occurs associated to shallow marine environments whereas their Recent counterparts are described on deep-marine corals. We interpret this as a consequence of a change in the environmental requirements of the coral/worm pair.

**Key words:** Scleractinia, Polychaeta, Eunicida, palaeoecology, bioerosion, commensalism, Pliocene, Mediterranean

Jordi Martinell [[jmartinell@ub.edu](mailto:jmartinell@ub.edu)] and Rosa Domènech [[rosa.domenech@ub.edu](mailto:rosa.domenech@ub.edu)], Dpt. d'Estratigrafia, Paleontologia i Geociències Marines, Facultat de Geologia, Universitat de Barcelona, c/ Martí i Franqués, s/n, E-08028 Barcelona, Spain.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

