

Mississippian colonial tabulate and rugose corals from the Flett Formation, Liard Basin, northwest Canada

Jerzy Fedorowski, E. Wayne Bamber, and Barry C. Richards *Acta Palaeontologica Polonica* 66 (3), 2021: 679-704 doi:https://doi.org/10.4202/app.00817.2020

Colonial Rugosa and Tabulata from the lower and middle Viséan Flett Formation of the Liard Basin (Canada) are described. *Michelinia expansa* is the first tabulate coral described from the Flett Formation. The new generic name *Cordilleria* is introduced for the North American Lithostrotionidae and its differences from the European *Siphonodendro*n are documented. Extremely wide variability in all characters and the small size of the studied collection allowed description of numerous Liard Basin specimens in open nomenclature only. Discussion of the most important characteristics of *Cordilleria* gen. nov. forms an introduction to the comprehensive revision of the North American Lithostrotionidae. The state of preservation of a colony identified as *Stelechophyllum* cf. *banffense* emphasizes the necessity of recognizing the diagenetic alterations of specimens prior to their taxonomic identification.

Key words: Rugosa, Tabulata, Favositina, Lithostrotionina, taxonomy, Carboniferous, Canada.

Jerzy Fedorowski [jerzy@amu.edu.pl], Institute of Geology, Adam Mickiewicz University, Bogumiła Krygowskiego 12, 61-680 Poznań, Poland.

E. Wayne Bamber [wayne.bamber@canada.ca] and Barry C. Richards [barry.richards@canada.ca], Geological Survey of Canada (Calgary) Natural Resources Canada, Government of Canada, 3303-33rd Street Northwest, Calgary, Alberta T2L 2A7, Canada.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see <u>creativecommons.org</u>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

