

## Evolution of the Jurassic and Cretaceous smooth-walled *Lenticulina* (Foraminiferida) of Poland

Bronisława Jendryka-Fuglewicz

*Acta Palaeontologica Polonica* 20 (2), 1975: 99-197

Three developmental phases of the *Lenticulina* have been distinguished, including (1) Lias, (2) Dogger - Lower Cretaceous and (3) Upper Cretaceous. The phylogenetic changes in the *Lenticulina* led to a decrease in the number of chambers in the outer whorl with a simultaneous increase in test size, in the degree of involution, and in the diameter of umbilical areas, as well as with a progressive development of the *Robulus* slit in aperture.

The decrease in the number of chambers in outer whorl occurs in independent ways, by: 1 - a decrease, in the geological time, of the number of chambers forming the test and 2 - a decrease in the number of chambers in outer whorl, with a simultaneous increase in the total number of chambers, in the length of spiral and number of whorls. Thirty-one species have been studied and morphotypes distinguished in polymorphic species. Simple statistical methods and microstructural studies were applied. The stratigraphic range of the following, so far confused, species: *Lenticulina gottingensis* (Bornemann), *L. muensteri* (Roemer) and *L. rotulata* Lamarck was established.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.