

Body size and diversity exemplified by three trilobite clades

Jerzy Trammer and Andrzej Kaim

Acta Palaeontologica Polonica 42 (1), 1997: 1-12

Cope's rule concerns only the radiation phase of a clade, overlooking the phase of the clade decline; thus it is incomplete. Changes of body size during the entire evolutionary history of a clade are exemplified here by three trilobite groups - Ptychopariina, Asaphina and Phacopida. Increasing diversity of the clade is associated with increase in maximum body size during the radiation phase, and decreasing diversity is generally associated with a decrease in maximum body size. Two basic patterns of the maximum body size changes are observed during the decline of the clade. The first one is characterized by a high correlation between diversity and the maximum body size, and indicative of species attrition that is nonselective with respect to the body size. The second one is characterized by a weak correlation between diversity and maximum body size, and typical of selective species attrition in relation to size.

Key words: body size, evolution, trends, trilobites

Jerzy Trammer, Instytut Geologii Podstawowej, Uniwersytet Warszawski, al. Żwirki i Wigury 93, PL-02-089 Warszawa, Poland; Andrzej Kaim, Instytut Paleobiologii PAN, ul. Twarda 51/53, PL-00-114 Warszawa, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.