

## Trace fossils from Late Carboniferous storm deposits, Upper Silesia Coal Basin, Poland

Arkadiusz Głuszek

*Acta Palaeontologica Polonica* 43 (3), 1998: 517-546

Eighteen ichnospecies (two new) referred to 17 ichnogenera (one new) as well as formally unclassified equilibrichnia, fugichnia and faecal pellets constitute a diverse and well-preserved invertebrate trace fossil suite present in storm deposits within the Franciszka X marine horizon of the Hruv Beds (Late Carboniferous Namurian A, Pendleian) as exposed in the Kozłowa Góra quarry, Upper Silesia, Poland. The new forms are *Fimbritubichnus biserialis* igen. et isp. n., which is interpreted as the work of a deposit-feeding bellerophont gastropod, and *Cylindrichnus candelabrus* isp. n. An emended diagnosis of the ichnospecies *Parahaentzschelinia ardeliais* given. Presence of the trace fossils marks a marine influence, so that they can be utilized to discern marine versus nonmarine deposition within the Paralic Series of the Upper Silesia Coal Basin in which other paleontological evidence is scanty or lacking.

**Key words:** Invertebrate trace fossils, storm deposits, Namurian A, Upper Silesia Coal Basin.

Arkadiusz Głuszek, Institute of Geological Sciences, Polish Academy of Sciences, ul. Senacka 1, PL-31-002 Kraków, Poland; the present adress: ul. Mickiewicza 19/14, PL-42-480 Poręba, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.