

Silurian retiolitid graptolites: Morphology and evolution

Denis E.B. Bates, Anna Kozłowska, and Alfred C. Lenz *Acta Palaeontologica Polonica* 50 (4), 2005: 705-720

Developmental mode and proximal structures are commonly accepted as the best for the recognition of high-level taxonomic categories within the Graptoloidea. The petalolithids and retiolitids are unique in possessing a virgellar ancora and in the latter, distal ancora development. The ancora structures are considered homologous, and the ancorate petalolithids are considered to be the direct ancestors to the retiolitids. The Retiolitidae are unique among the diplograptoids in possessing (1) outer, lateral, ancora sleeve walls (derived from distal extension of the ancora), and (2) a skeletal framework of bandaged lists between which are a succession of very thin and rarely preserved fusellar layers. Retiolitids possess different kinds of thecal profiles and two types of micro-ornamentation on the lists, and these have served to distinguish between the subfamilies Retiolitinae and Plectograptinae. Complete retiolitid morphological terminology is clarified and explained. Cladistic analysis of the retiolitids provides some measure of a better understanding of retiolitid evolution, but adds only modest support for the retention of the two subfamily categories.

Key words: Graptoloidea, Retiolitidae, Petalolithidae, ancora, cladistic analysis, Silurian.

Denis E.B. Bates [deb@aber.ac.uk], Institute of Geography and Earth Sciences, University of Wales, Aberystwyth Ceredigon SY23 3DB, UK; Anna Kozłowska[akd@twarda.pan.pl], Instytut Paleobiologii PAN, ul. Twarda 51/55, PL-00-818 Warszawa, Poland; Alfred C. Lenz [aclenz@uwo.ca], Department of Earth Sciences, University of Western Ontario, London, Ontario N6A 5B7, Canada.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see <u>creativecommons.org</u>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Full text (945.0 kB)