

## A stem-group frog from the Early Triassic of Poland

Susan E. Evans and Magdalena Borsuk-Białynicka

*Acta Palaeontologica Polonica* 43 (4), 1998: 573-580

Described on the basis of disarticulated postcranial bones (vertebrae, humeri, ilia) from karst deposits of Scythian age at the locality of Czatkowice in the Kraków Upland, Poland, *Czatkobatrachus polonicus* gen. et sp. n. is the first salientian known from the Triassic of the Northern part of Pangea. It may be only slightly younger (about 5 MA) than *Triadobatrachus massinoti* (Piveteau, 1936) from Madagascar, the only Early Triassic salientian known hitherto. *Czatkobatrachus* resembles *Triadobatrachus* but is more derived in some features of the vertebrae and elbow joint. It provides evidence of a global distribution of stem-frogs at the very beginning of the Mesozoic, and suggests that the origin of the group must be sought in the Permian.

**Key words:** Anura, Lissamphibia, Salientia, Triassic, Poland.

Susan E. Evans [[ucgasue@ucl.ac.uk](mailto:ucgasue@ucl.ac.uk)], Department of Anatomy and Developmental Biology, University College London, Gower Street, London, WC1E 6BT, England. Magdalena Borsuk-Białynicka [[borsuk.b@twarda.pan.pl](mailto:borsuk.b@twarda.pan.pl)], Instytut Paleobiologii PAN, ul. Twarda 51/55, PL-00-818 Warszawa, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.