

A new species of *Glyphea* (Decapoda: Palinura) from the La Meseta Formation (Eocene) of Seymour Island, Antarctica

Rodney M. Feldmann and Andrzej Gaździcki

Acta Palaeontologica Polonica 42 (3), 1997: 437-445

A new species of palinuran lobster, *Glyphea reticulata*, from the lowermost part of the Eocene La Meseta Formation on Seymour Island, Antarctica, represents one of the stratigraphically youngest species of *Glyphea*. The occurrence of the last vestiges of what was previously a cosmopolitan genus in a region dominated by Pacific Ocean faunal influences is significant because the sole extant species of the Glypheidae, *Neoglyphea inopinata* Forest & Saint Laurent, 1975, is known only from the west Pacific.

Key words: Decapoda, Glypheidae, paleobiogeography, evolution, Eocene, Antarctica.

Rodney M. Feldmann [rfeldman@kentvm.kent.edu], Department of Geology, Kent State University, Kent, Ohio, 44242, U.S.A. Andrzej Gaździcki [gazdzick@twarda.pan.pl], Instytut Paleobiologii PAN, ul. Twarda 51/55, PL-00-818 Warszawa, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.