



http://app.pan.pl/SOM/app59-Garilli_Parrinello_SOM.pdf

SUPPLEMENTARY ONLINE MATERIAL FOR

Taxonomy and palaeobiogeography of the Cenozoic Euro-Mediterranean rissoid gastropod *Galeodinopsis* and its relationship with close genera

Vittorio Garilli and Daniela Parrinello

Published in *Acta Palaeontologica Polonica* 2014 59 (2): 379–406.
<http://dx.doi.org/10.4202/app.2012.0044>

Supplementary Online Material

Table S1. Dimensions/main teleoconch sculpture parameters of *Galeodinopsis* spp. Abbreviations: AR, Agatino Reitano; AV, average value; Awp, number of axial ribs on the penultimate whorl; Coll., collection; σ , standard deviation; Ha, aperture height; Hlw, last whorl height; Ht, total shell height; IRSNB, Institut Royal des Sciences Naturelles de Belgique, Brussels; MF, Maurizio Forli; MNHU, Museum für Naturkunde der Humboldt Universität, Berlin; Sfw, number of spiral cords on the first whorl; Slw, number of spiral cords on the last whorl; SMF, Senckenberg Forschungsinstitut und Naturmuseum, Malakologie, Frankfurt am Main; Spw, number of spiral cords on the penultimate whorl; SS, Selected shell; UCBL, Université Claude Bernard Lyon 1; W, width.

Species	Specimen(s)/type of measurements	Ht (mm)	Hlw (mm)	Ha (mm)	W (mm)	Sfw	Spw	Slw	Awp
<i>Galeodinopsis biangulata</i> (Deshayes)	AV over 65-71 specimens	3.43	2.34	1.55	2.09	2	2	6.60	14.73
	♂ over 65-70 specimens	0.69	0.45	0.29	0.34	0	0	0.55	1.34
	SS, UCBL Deshayes coll., Oligocene, Paris Basin	4.95	3.15	1.95	2.50	2	2	6	14
	“	4.20	3.00	2.00	3.00	2	2	7	14
	SS, MNHU MB.Ga.7744, Oligocene, North Sea Basin,	4.30	2.60	1.90	2.70	2	2	6	15
	“	4.00	3.00	1.90	2.50	2	2	-	18
	“	3.40	2.50	1.60	2.10	2	2	7	16
<i>Galeodinopsis germanica</i> nov. sp.	SS, MF coll., Oligocene, North Sea Basin	3.20	2.40	1.50	2.00	2	2	7	15
	Holotype (SMF 335576)	2.55	1.55	1.10	1.50	4-5	6-7	14	18
	Paratype (SMF 336787)	2.30	1.50	1.05	1.40	4-5	6-7	14	17
	Paratype (SMF 336788)	2.00	1.35	0.90	1.17	4-5	5-6	11	20
	Paratype (SMF 336789)	1.72	0.95	0.70	1.11	4-5	5-6	13	19
	Paratype (SMF 336790)	1.90	1.20	0.80	1.15	4-5	5-6	14	20
	Paratype (SMF 336791a)	1.40	1.10	0.70	-	4-5	-	12	-
<i>Galeodinopsis semperi</i> (Wiechmann)	Paratype (SMF 336791b)	2.00	1.40	0.80	1.20	4-5	5-6	12	22
	IRSNB 3956, Nyst coll., Late Tongrian, Belgium	2.55	1.80	1.30	1.60	3	3-4	8	18
	Specimen MNHU MB.Ga.7745, Rupelian, Germany	3.52	2.36	1.57	2.16	4	4	10	24
	“	4.13	3.00	1.88	2.61	-	4	9	>20
	“	3.00	2.00	-	1.80	3-4	4	9	20
	Specimen in Tembrock (1964, pl. 7, fig. 6)	2.47	1.62	1.15	1.60	3	3-4	7	18
	Specimen in Tembrock (1964, pl. 7, fig. 7)	1.41	0.94	0.65	1.01	3	-	7	-
<i>Galeodinopsis tiberiana</i> (Coppi)	AV over more than 100 specimens	3.37	2.15	1.48	2.37	3	4.06	10.57	13.29
	♂ over more than 100 specimens	0.38	0.27	0.17	0.27	0	0.44	1.14	1.48
	SS, ex MF, MZDP Mal 2034 and 2035, Pliocene of N Italy	3.75	2.50	1.75	2.75	3	4	9	14
	“	3.75	2.25	1.65	2.25	3	4	10	13
	“	3.65	2.15	1.60	2.60	3	4	13	12
	“	3.65	2.25	1.55	2.70	3	4	11	9
	“	3.65	2.20	1.70	2.45	3	4	10	11
	“	3.60	2.10	1.55	2.60	3	4	9	13
	“	3.55	2.25	1.55	2.50	3	4	9	14
	“	3.50	2.15	1.50	2.75	3	4	8	14
	“	3.45	2.15	1.45	2.30	3	4	10	12
	“	3.40	2.35	1.45	2.35	3	4	9	10
	“	3.35	2.30	1.40	2.30	3	4	8	14
	“	3.30	2.10	1.35	2.25	3	4	9	12
	“	3.30	2.40	1.75	2.30	3	4	9	14
	“	3.30	2.45	1.75	2.45	3	4	10	14
	“	3.20	2.15	1.45	2.20	3	5	10	12
	“	3.20	2.10	1.35	2.15	3	4	11	12
	SS, ex AR, MZPD Mal 2034, Pliocene of Altavilla, S Italy	2.95	2.20	1.60	2.20	3	4	10	13
	“	2.80	2.00	1.50	2.00	3	4	11	13
“	2.50	1.80	1.30	1.75	3	4	11	12	
“	2.50	2.10	1.50	2.00	3	4	10	14	
<i>Galeodinopsis</i> cf. <i>tiberiana</i>	Specimen, ex AR, MZPD Mal 2033, Pliocene of Altavilla, S Italy	3.20	2.15	1.50	2.05	3	3	10	18
	“	2.93	2.00	1.35	1.85	3	5-6	10	20
	“	2.80	1.90	1.35	1.90	3	4	11	19
	“	2.57	1.80	1.25	1.65	3	4	9	18