

SUPPLEMENTARY ONLINE MATERIAL FOR

**Terrestrial-aquatic wood-inhabiting ascomycete *Potamomyces* from the
Miocene of Poland**

Grzegorz Worobiec, Elżbieta Worobiec, Przemysław Gedl, Jacek R. Kasiński,
Danuta Peryt, and Marek Widera

Published in *Acta Palaeontologica Polonica* 2022 67 (X): xxx-xxx.
<https://doi.org/10.4202/app.00976.2022>

Supplementary Online Material

Table 1. Modern and fossil record of ascospores considered as belonging to the *Potamomyces* (partly after Schlutz and Shumilovskikh 2013 and Nuñez Otaño et al. 2017)

Taxon name	Age	Location	Environment (source/substrate)	Climate	Reference
<i>Mediaverrunites fourierii</i> , <i>M. magnus</i> , <i>M. sp.</i>	early Miocene	Colombia	(sediment)	?	Elsik and Jarzen (2009)
<i>Mediaverrusporites sp.</i>	middle Miocene	western Niger Delta, Nigeria	freshwater to brackish (nearshore/coastal deltaic, swampy, mangrove sediment)	humid tropical	Durugbo and Olayiwola (2017)
<i>Mediaverrunites sp.</i>	middle/late Miocene	Arunachal Pradesh, India	freshwater to brackish (sediment)	humid tropical/subtropical	Paruya et al. (2017)
<i>Mediaverrunites mulleri</i> , <i>M. batii</i> , <i>M. pontidiensis</i>	late Miocene	Black Sea, Turkey	freshwater to marine (fluvio-deltaic to shallow marine sediment)	tropical to warm temperate	Sancay (2014)
<i>Mediaverrunites elsikii</i> , <i>M. mulleri</i>	Miocene	Mizoram, India	brackish (sediment)	tropical	Nandi and Sinha (2007)
<i>Mediaverrunites elsikii</i> , <i>M. mulleri</i>	Miocene	Solimões Formation, Amazon region, North	freshwater (lacustrine to fluvial)	humid tropical	Guimarães et al. (2013), da Silveira and

		Brazil	sediment)		Souza (2016)
<i>Mediaverrunites</i> sp.	Miocene	Barreiras Group, Amazon region, North Brazil	freshwater to marine (fluvio-deltaic to shallow marine sediment)	?	Arai et al. (1988)
<i>Mediaverrusporonites</i> sp.	Miocene	North of Chiapas, Mexico	(sediment)	probably tropical	Palacios-Chávez (1982)
<i>Potamomyces</i> sp. (<i>Potamomyces</i> sp. cf. <i>Mediaverrunites</i> sp. of Gemeraad 1979)	Miocene	Heath Formation, Tumbes Province, Peru	marine (sediment)	?	O'Keefe (2017)
<i>Mediaverrunites elsikii</i>	Miocene/Pliocene	Novo Remanso Formation, central Amazonia, Brazil	freshwater (meandering river sediment)	humid tropical	Guimarães et al. (2015)
<i>Mediaverrunites</i> sp.	Miocene/Pliocene	Siwalik strata, Churanthi River, India	brackish to shallow marine (sediment)	humid tropical to subtropical	More et al. (2016)
<i>Monoporisporites koenegii</i> (fig. 6S)	late Neogene	Mahuadanr valley, Jharkhand, India	(sediment)	humid tropical	Singh and Chauhan (2008), Singh and Prasad (2009)
<i>Mediaverrunites invaginatus</i>	late Neogene to early Pleistocene	offshore Louisiana Gulf of Mexico, USA	marine (sediment)	?	Elsik and Jarzen (2009)
Fungal spore (pl. 1: 2)	younger Cenozoic	cores from site 493 near Mexico	(sediment)	?	Fournier (1982)
<i>Potamomyces</i> sp.	late Quaternary	Rio Grande do Sul, southern Brazil	marine (coastal sediment)	subtropical	Roth and Lorscheitter (2016)
<i>Potamomyces</i> sp.	late Pleistocene	offshore northern Brazil	marine (offshore sediment)	tropical	Unpubl. data, Dupont et al. (2009) in: Schlütz and Shumilovskikh (2013)
<i>Mediaverrunites</i> sp.	late Pleistocene/Holocene	Laçador Palm Swamp, Buritizeiro, Minas Gerais, Brasil	swamp (sediment)	tropical	Cassino and Meyer (2013)
<i>Mediaverrunites</i> sp.	Holocene	New South Wales, Australia	(sediment)	subtropical	Elsik and Jarzen (2009)
<i>Mediaverrunites</i> sp.	Holocene	American Memorial Park, Saipan, Northern Marianas	brakish (mangrove sediment)	humid tropical	Jarzen and Dilcher (2006)
<i>Mediaverrunites</i> sp.	Holocene	Coastal Plain of Santa Catarina, southern Brazil	marine (nearshore/coastal sediment)	humid tropical	da Silva, and de Souza (2020)
<i>Mediaverrunites</i> sp.	Holocene	Mahwaqa Mountain, KwaZulu-Natal, South Africa	terrestrial (wetland sediment)	warm temperate, montane	Neumann et al. (2014)
<i>Potamomyces</i> sp.	Holocene	Yacata, Fiji Islands	freshwater (estuary/river sediment)	tropical	Macphail and Hope (2006)
<i>Potamomyces</i> sp.	Holocene	Eastern Drakensberg, South Africa	(sediment)	warm temperate	F. Neumann in: Schlütz and Shumilovskikh (2013)
Fungal ascospore (fig. 7BH)	Holocene	Wadhvana Lake, Gujarat, India	freshwater (fluvial sediment)	subtropical/tropical	Prasad et al. (2014)
Type JAM. 2053 (pl.6: 5)	Holocene	Jamaica	swamp (sediment)	tropical	Gemeraad (1979)
<i>Potamomyces palmarensis</i>	late Holocene	El Palmar National Park, Argentina	terrestrial (sediment)	humid sub-tropical	Nuñez Otaño et al. (2017)
<i>Potamomyces</i>	late Holocene	harbour of Elaia, ancient Pergamon, western Turkey	shallow marine (nearshore sediment)	subtropical	Shumilovskikh et al. (2016)
Ascospore (fig. 10i)	late Holocene	Cauca River, Colombia	freshwater (fluvio-palustrine sediment)	humid tropical	Serena et al. (2015)
<i>Mediaverrunites</i> sp.	Sub-recent	Sydney Cove (Warrane), New South Wales, Australia	terrestrial (archaeological layers)	subtropical	Macphail and Owen (2018)
<i>Mediaverrusporonites</i> sp.	Sub-recent	Parramatta, Australia	terrestrial (archaeological layers)	subtropical	Macphail (2006)
<i>Potamomyces</i>	Sub-recent	Yaque River valley, Dominican Republic	freshwater (fluvial sediment)	humid tropical	Hooghiemstra et al. (2018)
<i>Biconiosporella corniculata</i>	Recent	Anduki Beach, Brunei	brakish (driftwood from sandy sea shore)	tropical	Hyde (1988), Hyde (1995)
<i>Mediaverrunites</i> sp.	Recent	Burleson County, Texas, USA	terrestrial (humus-rich layer in hollow	warm temperate	Elsik and Jarzen (2009)

			tree)		
<i>Mediaverrunites</i> sp.	Recent	Luangwa valley, Zambia	freshwater (river sediment)	tropical	Jarzen and Elsik (1986)
<i>Potamomyces aquatica</i>	Recent	Tai Po Kau Forest Stream in Tai Po Kau Special Area, Hong Kong	freshwater (wood in stream)	subtropical	Ho et al. (2001)
<i>Potamomyces armatisporus</i>	Recent	Clohesy river, Australia	freshwater (wood in river)	humid tropical	Hyde (1995)
<i>Potamomyces armatisporus</i>	Recent	Rio Cuyabeno, Ecuador	freshwater (wood in stream)	tropical	Hyde (1995)
<i>Potamomyces armatisporus</i>	Recent	'Old Gombak', Malaysia	freshwater (wood)	tropical	Hyde (1995)
<i>Potamomyces armatisporus</i>	Recent	Lake Maicá, Santarém, Pará, Brazil	freshwater (wood in lake)	humid tropical	Canto et al. (2020)
<i>Potamomyces armatisporus</i> , <i>P. nepalensis</i>	Recent	Central Nepal	surface sample	subtropical	Schlütz and Shumilovskikh (2013)
Ascospore	Recent	Playa Medina, Venezuela	surface sample (coastal and mangrove)	humid tropical	Rull and Vegas-Vilarrubia (1999)
Fungal spore (pl. 1.:10)	Recent	Orinoco delta, Colombia	freshwater/brakish (deltaic, nearshore sediment)	humid tropical	Muller (1959)
<i>Striadiporites</i> sp. (fig. 3H)	Recent	Fazenda Urbano, Buritizeiro municipality, Minas Gerais, Brazil	swamp (sediment)	tropical	Lorente and Meyer (2010)

References

- Arai, M., Uesugui, N., Rossetti, D.D.F. and Góes, A.M. 1988. Considerações sobre a idade do Grupo Barreiras no nordeste do Estado do Pará. In: *Congresso Brasileiro de Geologia*, 35(1988): 738–752.
- Canto, E.S.M., Cortez, A.C.A., Monteiro, J.S., Barbosa, F.R., Zelski, S. and Souza, J.V.B.D. 2020. Composition and Diversity of Fungal Decomposers of Submerged Wood in Two Lakes in the Brazilian Amazon State of Para. *International Journal of Microbiology*: 2020.
- Cassino, R.F., and Meyer, K.E.B. 2013. Reconstituição paleoambiental do Chapadão dos Gerais (Quaternário tardio) a partir da análise palinológica da Vereda Laçador, Minas Gerais, Brasil. *Revista Brasileira de Paleontologia*, 16(1): 127–146.
- da Silva, W.G., and de Souza, P.A. 2020. Cistos de dinoflagelados do Holoceno da planície costeira De Santa Catarina (Poço PSC-03): descrições taxonômicas e implicações paleoambientais. *Geociências (São Paulo)*, 38(3): 795–812.
- da Silveira, R.R., & Souza, P.A. 2016. Palinologia (esporos de fungos e pteridófitas, grãos de pólen de gimnopermas, cistos de algas e escolocodonte) das formações Solimões e Içá (Neogeno e Pleistoceno, Bacia do Solimões), Amazonas, Brasil. *Pesquisas em Geociências*, 43(1): 17–39.
- Dupont, L., Schlütz, F., Ewah, C., Jennerjahn, T.C., Paul, A. and Behling, H. 2009. Two-step vegetation response to enhanced precipitation in Northeast Brazil during Heinrich event 1. *Global Change Biology*, 10: 1647–1660.
- Durugbo, E.U. and Olayiwola, M.A. 2017. Palynological dating and palaeoenvironments of the M1 well, Middle Miocene, Niger Delta, Nigeria. *Palaeontologia africana*, 52: 46–57.
- Elsik, W.C., Jarzen, D.M. 2009. New species of the late Cenozoic fungal form-genus *Mediaverrunites* Jarzen und Elsik 1986 ex Nandi und Sinha 2007. *Palynology*, 33(2): 99–104.

- Fournier, G.R. 1982. Palynostratigraphic analysis of cores from site 493, Deep Sea Drilling Project leg 66. *Initial Reports of the Deep Sea Drilling Project*, 66: 661–670.
- Germeraad, J.H. 1979. Fossil remains of fungi, algae and other organisms from Jamaica. *Scripta Geologica*, 52: 1–39.
- Guimarães, J.T.F., Nogueira, A.C.R., Da Silva Jr, J.B.C., Soares, J.L. and Silveira, R. 2013. Fossil fungi from Miocene sedimentary rocks of the central and coastal Amazon region, North Brazil. *Journal of Paleontology*, 87(3): 484–492.
- Guimarães, J.T.F., Nogueira, A.C.R., Da Silva Jr, J.B.C., Soares, J.L., Alves, R., Kern, A.K. 2015. Palynology of the Middle Miocene-Pliocene Novo Remanso Formation, Central Amazonia, Brasil. *Ameghiniana*, 52(1): 107–134.
- Ho, W.H., Hyde, K.D. and Hodgkiss, I.J. 2001. Fungal communities on submerged wood from streams in Brunei, Hong Kong, and Malaysia. *Mycological Research*, 105(12): 1492–1501.
- Hooghiemstra, H., Olijhoek, T., Hoogland, M., Prins, M., van Geel, B., Donders, T., Gosling, W. and Hofman, C. 2018. Columbus' environmental impact in the New World: Land use change in the Yaque River valley, Dominican Republic. *The Holocene*, 28(11): 1818–1835.
- Hyde, K.D. 1988. Studies on the tropical marine fungi of Brunei. *Botanical Journal of the Linnean Society*, 98(2): 135–151.
- Hyde, K.D. 1995. Tropical Australasian fungi. VII. New genera and species of ascomycetes. *Nova Hedwigia*, 61(1-2): 119–140.
- Jarzen, D.M. and Dilcher, D.L. 2009. Palynological assessment of Holocene mangrove vegetation at the American Memorial Park, Saipan, Northern Mariana Islands. *Grana*, 48(2): 136–146.
- Jarzen, D.M., Elsik, W.C. 1986. Fungal palynomorphs recovered from recent river deposits, Luangwa Valley, Zambia. *Palynology*, 10: 35–60.
- Lorente, F.L. and Meyer, K.E.B. 2010. Palinomorfos da vereda da Fazenda Urbano, município de Buritizeiro, Minas Gerais, Brasil. *Iheringia, Série Botânica*, 65(2): 133–169.
- Macphail, M.K. 2006. Pollen analysis of soil samples, 109-113 George Street, Parramatta. In: M. Casey and T. Lowe (eds), *Archaeological Investigation Report Section 8.4 Pollen Report vol. 2*: 1–60. <http://www.caseyandlowe.com.au/pdf/109/v2section84.pdf>.
- Macphail, M. and Hope, G. 2006. Natural Histories - An Illustrated Guide to Fossil Pollen and Spores Preserved in Swamps and Mires on the Southern Highlands of NSW. The Australian National University, Canberra.
- Macphail, M. and Owen, T. 2018. What was growing along the Tank Stream Valley, Sydney Cove, in 1788?. *Australasian Historical Archaeology*, 36: 16–28.
- More, S., Paruya, D.K., Taral, S., Chakraborty, T. and Bera S. 2016. Depositional Environment of Mio-Pliocene Siwalik Sedimentary Strata from the Darjeeling Himalayan Foothills, India: A Palynological Approach. *PLoS ONE* 11(3): e0150168.
- Muller, J. 1959. Palynology of Recent Orinoco delta and shelf sediments; reports of the Orinoco shelf expedition, volume 5. *Micropaleontology*, 5(1): 1–32.
- Nandi, B. and Sinha, A. 2007. Validation of the Miocene fungal spore *Mediaverrunites* from Mizoram, India. *Palynology*, 31(1): 95–100.

- Neumann, F.H., Botha, G.A. and Scott, L. 2014. 18,000 years of grassland evolution in the summer rainfall region of South Africa: evidence from Mahwaqa Mountain, KwaZulu-Natal. *Vegetation History and Archaeobotany*, 23(6): 665–681.
- Nuñez Otaño, N., di Pasquo, M. and Bianchinotti, M.V. 2017. The occurrence of *Potamomyces palmarensis* sp. nov. in the Late Holocene of El Palmar National Park (Colón, Entre Ríos, Argentina) and transfer of fossil species of *Mediaverrunites* to *Potamomyces*. *Palynology*, 41(2): 267–277.
- O'Keefe, J.M.K. 2017. Fungal palynomorphs from the Miocene Heath Formation, Tumbes Province, Perú. *Palynology*, 41: 309–326.
- Palacios-Chávez, R. 1982. Las fungosporas del Mioceno inferior y medio del norte de Chiapas (Mexico). *Scientia Fungorum*, 2(17): 33–42.
- Paruya, D.K., Ghosh R., Biswas, O., Bera, M. and Bera, S. 2017. Dispersed fungal remains from the Neogene Siwalik forest of sub-Himalayan Arunachal Pradesh, India and their palaeoenvironmental indicative values. *Journal of Mycopathological Research*, 55(3): 303–307.
- Prasad, V., Farooqui, A., Sharma, A., Phartiyal, B., Chakraborty, S., Bhandari, S., Raj, R. and Singh, A. 2014. Mid–late Holocene monsoonal variations from mainland Gujarat, India: A multi-proxy study for evaluating climate culture relationship. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 397: 38–51.
- Roth, L. and Lorscheitter, M.L. 2016. Fungi, algae, and other palynomorphs in sedimentary profiles collected from two forests in the northernmost coastal plain from Rio Grande do Sul, southern Brazil. *Brazilian Journal of Botany*, 39(4): 1135–1143.
- Rull, V. and Vegas-Vilarrubia, T. 1999. Surface palynology of a small coastal basin from Venezuela and potential paleoecological applications. *Micropaleontology*, 45: 365–393.
- Sancay, R.H. 2014. The occurrence of *Mediaverrunites* in the Upper Miocene of the Black Sea, Turkey. *Palynology*, 38: 28–37.
- Schlütz, F. and Shumilovskikh, L.S. 2013. On the relation of *Potamomyces armatisporus* to the fossil form-type *Mediaverrunites* and its taxonomical and ecological implications. *Fungal Ecology*, 6(4): 309–315.
- Serna, Y., Velez, M.I., and Escobar, J. 2015. Microscopic organic matter particles in late Holocene riparian sediments near the Cauca River, Colombia. *Journal of paleolimnology*, 54(4): 325–344.
- Shumilovskikh, L.S., Seeliger, M., Feuser, S., Novenko, E., Schlütz, F., Pint, A., Pirson, F. and Brückner, H. 2016. The harbour of Elaia: A palynological archive for human environmental interactions during the last 7500 years. *Quaternary Science Reviews*, 149: 167–187.
- Singh, S.K. and Chauhan, M.S. 2008. Fungal remains from the Neogene sediments of Mahuadanr Valley, Latehar District, Jharkhand, India and their palaeoclimatic significance. *Journal of the Palaeontological Society of India*, 53(1): 73–81.
- Singh, S.K., and Prasad, M. 2009. Floral Diversity and Climate During Late Tertiary Period in Mahuadanr Valley, Jharkhand, India. *Phytomorphology*, 59: 19–28.