

A new Cambrian catillicephalid trilobite from the Shallow Bay Formation of western Newfoundland, Canada

Stephen R. Westrop and Alyce A. Dengler *Acta Palaeontologica Polonica* 67 (1), 2022: 27-33 doi:https://doi.org/10.4202/app.00922.2021

Species of *Catillicephala* are known from sites around the mid-Cambrian margin of Laurentian North America, including Vermont, Quebec, Newfoundland and North Greenland. *Catillicephala cifellii* sp. nov. is from the Downes Point Member of the Shallow Bay Formation (Cow Head Group) in western Newfoundland. It occurs in three shelf margin-derived boulders in debris flow conglomerates that accumulated in a continental slope setting. The associated trilobites and agnostoid arthropods, including *Ptychagnostus aculeatus* and *Megagnostus glandiformis*, indicate a correlation with the *Lejopyge laevigata* Zone. As such, *C. cifellii* is among the oldest representatives of the genus, and is early Guzhangian in age.

Key words: Trilobita, *Catillicephala*, Cambrian, Laurentia, North America.

Stephen R. Westrop [swestrop@ou.edu], Oklahoma Museum of Natural History and School of Geosciences, University of Oklahoma, Norman, OK 73072, USA. Correspondence address: 103 Melita Street, Ingersoll, Ontario N5C 3W3, Canada. Alyce A. Dengler [sandydengler2@yahoo.com], Oklahoma Museum of Natural History, University of Oklahoma, Norman, OK 73072, USA.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see <u>creativecommons.org</u>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.