

Photography in the ultraviolet and visible violet spectra: Unravelling methods and applications in palaeontology

Gaia Crippa and Stefano Masini

Acta Palaeontologica Polonica 67 (3), 2022: 685-702 doi:<https://doi.org/10.4202/app.00948.2021>

We have tested different preparation and photographic methods to define a protocol for UV analysis of fossil specimens. We also have explored its main applications while analysing specimens from different stratigraphic contexts, of different biomineralogical composition, and belonging to different fossil groups (including invertebrates and vertebrates). We have photographed specimens using a camera equipped with appropriate lens and filters both in visible light and with flashlights at two wavelengths: the 365 nm UV light and the 440 nm visible violet spectrum, the latter here tested for the first time. Our results indicate that bleach treatment is not recommended for calcite-shelled brachiopods, while it is suggested for aragonite-shelled molluscs. We show that photography in the ultraviolet and visible violet spectra are useful tools enhancing the recognition of morphological characters and colour patterns and allowing to distinguish soft-bodied fossils from the matrix. Also, it allows to discern specimen areas embedded in the sediment from those exposed to sunlight, which is helpful to reconstruct the conditions experienced by fossils. However, the mineralogy of the biomineral affects UV responses, as morphological characters of calcite shells are better emphasized with the 440 nm wavelength (visible violet spectrum), whereas those of aragonite, bioapatite and phosphatized specimens with the 365 nm (ultraviolet spectrum); also, shell microstructures with their different crystal arrangement and elemental incorporation may cause different reactions, whereas the stratigraphic context affects specimen preservation influencing pigment preservation. We thus provide a protocol for photography in the ultraviolet and visible violet spectra and show that this technique has a high potential in palaeontology, having no limitations for its application in invertebrate or vertebrate specimens.

Key words: Biominerals, fossil specimens, shell colour pattern, UV photography, visible violet spectrum.

Gaia Crippa [gaia.crippa@unimi.it] and Stefano Masini [stefano.masini92@gmail.com], Dipartimento di Scienze della Terra “A. Desio”, Università degli Studi di Milano, via Mangiagalli 34, Milano, 20133, Italy.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(1,220.4 kB\)](#) |

 [Supplementary file \(1,353.0 kB\)](#)