

## Triassic coleoid beaks and other structures from the Calcareous Alps revisited

Larisa A. Doguzhaeva, Herbert Summesberger, Franz Brandstaetter, Daniela Gruber, and Andrea Tintori  
*Acta Palaeontologica Polonica* 67 (3), 2022: 655-666 doi:<https://doi.org/10.4202/app.00953.2021>

We performed comprehensive study of seven Carnian, Late Triassic specimens of a coleoid cephalopod *Phragmoteuthis bisinuata*, on which Suess based his hypothesis on “beaks of *P. bisinuata*”. Using SEM/EDS, we found that “beaks of *P. bisinuata*” consist of a micro-granular carbonized matrix containing ~4–30 µm diameter and ~50–200 µm visible length, dense calcified bone-like micro-structures. This strongly suggests that these objects are vertebrate bone-inducing cartilages in which the matrix was post-mortem reworked by carbon-accumulating bacteria and substituted by nano-particles of carbon accumulated in micro-granules. Hence, the presumed “beaks of *P. bisinuata*” are cartilaginous remains of a prey, presumably juvenile fish. This data dismissed the entire hypothesis of Suess. A small spatula-shape plate with a rachis-like process in an association with 10 or so imprints around (arm crown), found in front of a proostracum of *P. bisinuata* evidences an unknown Late Triassic juvenile teuthid which possessed a gladius resembling that of the early Permian *Glochinomorpha stifeli*. It inhabited the open sea area of the northwestern Tethys Ocean, and was, along with juvenile fishes, in the diet of *P. bisinuata*. The first identified Anisian (Middle Triassic) coleoid beak is represented by an isolated specimen from the Gardena Valley, NE Italy. It has a typical composition and morphology of coleoid upper beak: chitinous, wide-oval lateral walls, short wings, and pointed hook-like rostrum. This suggests similar upper beak structure in the Carnian *P. bisinuata* in which the lower beaks were apparently similar to that of the co-occurring *Lunzoteuthis schindelbergensis* and had a widely open outer lamella with posteriorly elongated paired wings joined into a pointed rostrum in the anterior portion.

**Key words:** Cephalopoda, Coleoidea, *Phragmoteuthis*, beaks, vertebrate and invertebrate prey, Triassic, Alps.

Larisa A. Doguzhaeva [[larisa.doguzhaeva@gmail.com](mailto:larisa.doguzhaeva@gmail.com)],  
Department of Palaeobiology, Swedish Museum of Natural  
History, P.O. Box 50007, Stockholm SE-104 05, Sweden.  
Herbert Summesberger [[herbert.summesberger@nhm-wien.ac.at](mailto:herbert.summesberger@nhm-wien.ac.at)],  
Department of Geology and Palaeontology, Museum of Natural  
History, Vienna, Burgring 7, Austria. Franz Brandstaetter [[franz.brandstaetter@nhm-wien.ac.at](mailto:franz.brandstaetter@nhm-wien.ac.at)]  
, Mineralogical Department, Museum of Natural History, Vienna, Austria;

Vienna, Burgring 7, Austria. Daniela Gruber [[daniela.gruber@univie.ac.at](mailto:daniela.gruber@univie.ac.at)],  
Core Facility of Cell Imaging and Ultrastructure Research, Life Sciences  
Faculty, University of Vienna, 1090 Wien Althanstraße 14, Austria. Andrea  
Tintori [[paleo.tintori@outlook.it](mailto:paleo.tintori@outlook.it)], Triassica, Institute for Triassic Lagerstätte, 23828 Periedo (LC), Via al  
Verder 6, Italy.

This is an open-access article distributed under the terms of the Creative Commons  
Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use,  
distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(3,462.4 kB\)](#)