

## A new species of mixosaurid ichthyosaur from the Middle Triassic of Luxi County, Yunnan Province, South China

Ye-Wei Fang, Andrzej S. Wolniewicz, and Jun Liu

*Acta Palaeontologica Polonica* 69 (2), 2024: 263-280 doi:<https://doi.org/10.4202/app.01133.2024>

Ichthyosaurs, an iconic lineage of Mesozoic marine reptiles, were an important component of recovering ecosystems after the Permo-Triassic Mass Extinction event. Mixosauridae, a clade of small, early-diverging ichthyosaurs, were of particular significance for this process, being abundant predators in Middle Triassic shallow seas. Despite the abundance of well-preserved mixosaurid specimens from South China, *Mixosaurus panxianensis* remains the only comprehensively described species, hindering our understanding of the variability, taxonomy and diversity of mixosaurids from this region. Here, we report a new species of *Mixosaurus*, *Mixosaurus luxiensis*, from Luxi County, Yunnan Province, South China. The wider postorbital skull portion differentiates the new species from *Mixosaurus cornalianus* and *Mixosaurus kuhnschnyderi* from central Europe. The non-durophagous dentition, composed of tiny piercing mesial teeth and robust but pointed distal teeth, resembles the dentition of *M. cornalianus*. However, the distal teeth of *M. luxiensis* sp. nov. are twice the size of the mesial ones, in contrast to *M. cornalianus*, in which the mesial and distal teeth are approximately equal in size. The forelimb exhibits a unique morphology, including a proportionally narrow radius, the presence of a peripheral notch on the ulna, and a large metacarpal V. A preliminary phylogenetic analysis suggests a close affinity of the new taxon with *M. cornalianus* from Western Tethys. Our study introduces important, new anatomical information on *Mixosaurus* from South China, useful for future studies of mixosaurid diversity.

**Key words:** Ichthyosauria, Mixosauridae, marine reptile, phylogeny, biotic recovery, Guanling Formation, Anisian, Triassic.


Ye-Wei Fang [[paleofang@gmail.com](mailto:paleofang@gmail.com); ORCID: <https://orcid.org/0000-0002-8150-6331> ]

and Jun Liu [[junliu@hfut.edu.cn](mailto:junliu@hfut.edu.cn); ORCID: <https://orcid.org/0000-0001-7859-5209> ]

(corresponding author), School of Resources and Environmental Engineering, Hefei University of Technology, 193 Tunxi Road, Hefei 230009, Anhui, China. Andrzej S.

Wolniewicz [[asw49@cam.ac.uk](mailto:asw49@cam.ac.uk); ORCID: <https://orcid.org/0000-0002-6336-8916> ], School of Resources and Environmental Engineering, Hefei University of Technology, 193 Tunxi Road, Hefei 230009, Anhui, China; Department of Earth Sciences, University of Cambridge, Downing Street, Cambridge, CB2 3EQ, UK; Institute of Paleobiology, Polish Academy of Sciences, Twarda 51/55, 00-818 Warsaw, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(4,154.0 kB\)](#) |

 [Supplementary file \(518.6 kB\)](#)