

A Late Jurassic deep-bodied actinopterygian fish from Antarctica

Soledad Gouiric-Cavalli, Ari Iglesias, Bárbara Cariglino, and Marcelo A. Reguero
Acta Palaeontologica Polonica 69 (3), 2024: 467-483 doi:10.4202/app.01158.2024

Mesozoic deep-bodied actinopterygians are of interest given, among others, the various modes of feeding exhibited by these fishes. Regrettably, most of their fossil record is restricted to a limited number of localities in Europe. During the Late Jurassic fragmentation of Pangaea, the exchange of fauna between the European Tethys and the paleo-Pacific (southwestern of the South American region) was possible via the marine pathways connecting these areas. This exchange led to the speciation of fish taxa, particularly in the Southern Hemisphere. Although new species are continuously being discovered, our understanding of the Late Jurassic marine ichthyofaunas of the Southern Hemisphere remains limited. The Mesozoic ichthyofaunas of the Antarctic seas are rich but relatively poorly known presenting a significant opportunity for further research. Noteworthy, previous reports have documented the presence of actinopterygians in the Late Jurassic of the Antarctic Peninsula. Since 2016, a team of researchers from Argentina has been exploring the Upper Jurassic–Lower Cretaceous Ameghino (= Nordenskjöld) Formation outcrops, resulting in the discovery of a large collection of actinopterygian fishes. Nevertheless, most of the material remains undescribed. In this article, we report the discovery of a deep-bodied actinopterygian found at the Longing Gap, the type locality of the Ameghino Formation in the Antarctic Peninsula. The study of these newly collected materials allows for their taxonomic assignment to *Ameghinichthys antarcticus*, a taxon previously described for the locality but based on isolated and fragmentary material. Additionally, this study confirms that *A. antarcticus* belongs to Dapediiformes. *Ameghinichthys antarcticus* is among the youngest records of Dapediiformes and represents the southernmost record of the group worldwide.

Key words: Actinopterygia, Dapediiformes, Dapediidae, ichthyofaunas, Mesozoic, Gondwana.

Soledad Gouiric-Cavalli [sgouiric@fcnym.unlp.edu.ar; ORCID: <https://orcid.org/0000-0003-2026-5973>], División Paleontología de Vertebrados, Museo de La Plata, Universidad Nacional de La Plata (UNLP). Paseo del Bosque s/n, 1900 La Plata, Buenos Aires, Argentina. Consejo Nacional de Investigaciones Científicas y Tecnológicas (CONICET), Godoy Cruz 2290, C1425FQB Ciudad Autónoma de Buenos Aires, Argentina. Ari Iglesias [ari_iglesias@yahoo.com.ar; ORCID: <https://orcid.org/0000-0002-9098-8758>], Instituto de Investigaciones en

Biodiversidad y Medioambiente (CONICET-UNCO). Pasaje Gutiérrez 1415, R8400GAO, San Carlos de Bariloche, Río Negro, Argentina. Bárbara Cariglino[barichi10@gmail.com ; ORCID: <https://orcid.org/0000-0002-4346-3502>], División Paleobotánica y Paleopalinología, Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”, Av. Ángel Gallardo 470, C1405DJR, Ciudad Autónoma de Buenos Aires, Argentina. Marcelo A. Reguero [regui@fcnym.unlp.edu.ar ; ORCID: <https://orcid.org/0000-0003-0875-8484>], Instituto Antártico Argentino, Dirección Nacional del Antártico. Balcarce 295, C1064AAF Ciudad Autónoma de Buenos Aires, Argentina and División Paleontología de Vertebrados, Museo de La Plata, Universidad Nacional de La Plata (UNLP). Paseo del Bosque s/n, 1900 La Plata, Buenos Aires, Argentina. Consejo Nacional de Investigaciones Científicas y Tecnológicas (CONICET), Godoy Cruz 2290, C1425FQB Ciudad Autónoma de Buenos Aires, Argentina.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(2,230.7 kB\)](#)