

Life strategies and function of dissepiments in rugose coral *Catactotoechus* instabilis from the Lower Devonian of Morocco

Błażej Berkowski

Acta Palaeontologica Polonica 57 (2), 2012: 391-400 doi: http://dx.doi.org/10.4202/app.2010.0048

This study focuses on the life strategies of small, dissepimented rugose coral *Catactotoechus instabilis* (representative of *Cyathaxonia* fauna) from the Emsian argillaceous deposits of mud mounds of Hamar Laghdad (Anti–Atlas, Morocco).

Numerous constrictions and rejuvenescence phenomena as well as frequent deflections of growth directions among the studied specimens suggest unfavourable bottom conditions resulted from sliding down of the soft sediment on the mound slopes. Dissepimental structures observed on well–preserved calices and thin sections played an important role in the life of the coral, supporting their successful recovery after temporary burial within unstable soft sediment. The development of lonsdaleoid dissepiments, apart from being biologically controlled, was also strongly influenced by environmental factors. Such modifications in lonsdaleoid dissepiments growth were observed in phases of constrictions, rejuvenescence and deflections of growth, when their development was significantly increased in comparison to phases of their stable growth. Dissepiment morphology suggests that the process of formation of lonsdaleoid dissepiments in *Catactotoechus instabilis* is consistent with the hydraulic model.

Key words: Rugosa, corals, dissepiments, life strategies, Devonian, Hamar Laghdad, Morocco.

Błażej Berkowski [bbrk@amu.edu.pl], Institute of Geology, Adam Mickiewicz University, Maków Polnych 16, 60-606 Poznań, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see <u>creativecommons.org</u>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

