

Osteology of the Late Cretaceous alvarezsauroid *Linhenykus monodactylus* from China and comments on alvarezsauroid biogeography

Xing Xu, Paul Upchurch, Qingyu Ma, Michael Pittman, Jonah Choiniere, Corwin Sullivan, David W.E. Hone, Qingwei Tan, Lin Tan, Dong Xiao, and Fenglu Han

Acta Palaeontologica Polonica 58 (1), 2013: 25-46 doi: <http://dx.doi.org/10.4202/app.2011.0083>


The alvarezsauroid theropod *Linhenykus monodactylus* from the Upper Cretaceous of Inner Mongolia, China is the first known monodactyl non-avian dinosaur, providing important information on the complex patterns of manual evolution seen in alvarezsauroids. Here we provide a detailed description of the osteology of this taxon. *Linhenykus* shows a number of features that are transitional between parvicursorine and non-parvicursorine alvarezsauroids, but detailed comparisons also reveal that some characters had a more complex distribution. We also use event-based tree-fitting to perform a quantitative analysis of alvarezsauroid biogeography incorporating several recently discovered taxa. The results suggest that there is no statistical support for previous biogeographic hypotheses that favour pure vicariance or pure dispersal scenarios as explanations for the distributions of alvarezsauroids across South America, North America and Asia. Instead, statistically significant biogeographic reconstructions suggest a dominant role for sympatric (or “within area”) events, combined with a mix of vicariance, dispersal and regional extinction. At present the alvarezsauroid data set is too small to completely resolve the biogeographic history of this group: future studies will need to create larger data sets that encompass additional clades.

Key words: Dinosauria, Theropoda, Parvicursorinae, alvarezsauroid biogeography, Treefitter, dispersal, vicariance, sympatry, Cretaceous, Wulansuhai Formation, Inner Mongolia, China.

Xing Xu [xingxu@vip.sina.com], Qingyu Ma [maqingyu.ivpp@gmail.com], Corwin Sullivan [csullivan@ivpp.ac.cn], David W.E. Hone [dwe_hone@yahoo.com], and Fenglu Han [hfl0501@gmail.com], Key Laboratory of Evolutionary Systematics of Vertebrates, Institute of Vertebrate Paleontology & Paleoanthropology, Chinese Academy of Sciences, 142 Xiwai Street, Beijing 100044, China; Paul Upchurch [p.upchurch@ucl.ac.uk] and Michael Pittman [mdpittman@hotmail.com], Department of Earth Sciences, University College London, Gower Street, London, WC1E 6BT, UK; Jonah Choiniere [jonah.choiniere@gmail.com], Department of Biological Sciences, George Washington University, 2023 G Street NW, Washington, DC 20052, USA; Qingwei Tan [firsttan@sina.com] and Lin Tan, Long Hao Institute of Geology and Paleontology, Hohhot, Nei Mongol 010010, China; Dong Xiao, Department of Land and

Resources, Linhe, Nei Mongol 015000, China.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(1,005.6 kB\)](#) |

 [Supplementary file \(22.6 kB\)](#)