

## Some terebratulid populations from the Lower Kimmeridgian of Poland and their relations to the biotic environment

Jerzy Dzik

*Acta Palaeontologica Polonica* 24 (4), 1979: 473-492

Populations of *Epithyris* '*subsella*' in different Lower Kimmeridgian fossil assemblages show different size-frequency distributions although they exhibit a similar shape of survivorship curves. A population with large-sized adult specimens of *E.* '*subsella*' was characteristic for a community in which most abundant were bivalves: *Isognomon subplana*, *Lopha gregarea* and *Trichites saussurei*. A population with small adult specimens was characteristic for a community with abundant dasyclad alga *Goniolina geometrica*. Here, among shelled suspension-feeders the most important producers were brachiopods *E.* '*subsella*', *Zeilleria humeralis*, and *Septaliphoria pinguis*. Thalli of *G. geometrica* were the substrate for most of the sedentary organisms in this community. The estimation of biomass and productivity of particular species in fossil communities is discussed.

**Key words:** Brachiopods, fossil assemblages, Upper Jurassic, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(1,085.4 kB\)](#)