

Mid-Cretaceous microbiostratigraphy and foraminifers of the NE margins of the Świętokrzyskie (Holy Cross) Mts., Poland

Danuta Peryt

Acta Palaeontologica Polonica 28 (3-4), 1983: 417-466

Mid-Cretaceous foraminifers of the northeastern Mesozoic margin of the Holy Cross Mts. have been described. Planktic foraminifer standard zones - the *Rotalipora appenninnica*, *Rotalipora brotzeni*, *Rotalipora cushmani*, *Praeglobotruncana helvetica* - have been distinguished in the profile of Ożarów. A local benthic zonation, with the *Lingulogavelinella frankei*, *Gavelinella schloenbachi*, *Lingulogavelinella spinosa*, *Lingulogavelinella formosa*, *Gavelinella baltica* and *Lingulogavelinella globosa* zones, has been established in this profile. Due to the strong stratigraphic condensation and mixed fauna, no boundaries between the zones mentioned could be determined in the Rachów anticline.

Key words: Foraminiferida, Albian, Cenomanian, Turonian, stratigraphy, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(3,194.5 kB\)](#)