

## Middle Triassic Scleractinia from the Cracow-Silesia region, Poland

Elżbieta Morycowa

*Acta Palaeontologica Polonica* 33 (2), 1988: 91-121

Scleractinian coral buildups from the Karchowice Beds (uppermost part of the Lower Muschelkalk) and from the Diplopora Dolomite (Middle Muschelkalk) of the Cracow-Silesia region were examined. Described corals comprise: three species known from literature-*Coelocenia* cf. *decipiens* (Laube), *Pamiroseris silesiaca* (Beyrich), *Eckastraea prisca* (Weissermel); four new species -*Volzeia szulci*, *Morycastraea opoliensis*, *Silesiastraea weissermeli*, *Chevalieria tenuiseptata*; and three determined as *Stylophyllopsis* sp., *Cyathocoenia* sp. and *Pamiroseris* sp. New genera, *Eckastraea* and *Sileriastraea*, are established. The most frequent component of coral communities appears *Pamiroseris silesiaca*.

**Key words:** Scleractinia, taxonomy, morphology, paleoecology, Middle Triassic, southern Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



[Full text \(2,323.0 kB\)](#)