

## Devonian phoebodont shark teeth

Michał Ginter and Alexander Ivanov

*Acta Palaeontologica Polonica* 37 (1), 1992: 55-75

Shark teeth of the phoebodont type are the most common and diverse group of Upper Devonian ichthyoliths in the pelagic facies of the Holy Cross Mountains (Poland), South Urals and Timan (Russia). They were also found in the Givetian of Kuznetsk Basin (western Siberia). The morphology and function of tooth apparatus of *Phoebodus* was possibly similar to that of the recent shark *Chlamydoselachus anguineus*. A significant loss of diversity and relative productivity has been observed among the phoebodonts in the earliest Famennian. A new genus, *Omalodus* gen. n., and three new species of *Phoebodus*, *Ph. bifurcatus* sp. n., *Ph. fastigatus* sp. n. and *Ph. turnerae* sp. n. are proposed.

**Key words:** sharks, phoebodonts, teeth, morphology, biostratigraphy, Devonian, Poland, Russia.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



[Full text \(1,233.8 kB\)](#)