

The quadrate of oviraptorid dinosaurs

Teresa Maryńska and Halszka Osmólska

Acta Palaeontologica Polonica 42 (3), 1997: 361-371

The quadrate of oviraptorid dinosaurs is strongly pneumatized and differs from the quadrates of other non-avian theropods by: (1) two separate facets on the otic process for contacts with the squamosal and braincase; (2) the articular surface for the pterygoid extended to the articular surface of the medial mandibular condyle; (3) the mandibular process provided laterally with a quadratojugal process bearing the quadratojugal cotyla. In the above characters the oviraptorid quadrate resembles those in most ornithothoracine birds, but, contrary to the streptostylic quadrate of birds, the oviraptorid quadrate is monimostylic.

Key words: Dinosauria, Oviraptoridae, quadrate.

Teresa Maryńska, Muzeum Ziemi, PAN, al. Na Skarpie 20/26, PL-00-488

Warszawa, Poland. Halszka Osmólska [osm@twarda.pan.pl], Instytut Paleobiologii PAN, ul. Twarda 51/55, PL-00-818 Warszawa, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(1,870.9 kB\)](#)