

Body mass estimation and locomotion of the Miocene pelecaniform bird *Macranhinga*

Jorge Ignacio Noriega

Acta Palaeontologica Polonica 46 (2), 2001: 247-260

The fossil darter *Macranhinga paranensis* (Aves: Pelecaniformes) from the late Miocene of Argentina is the largest of all known extinct or living Anhingidae. Its body mass is estimated at a mean value of 5.4 kg by using a scaling model derived from the logarithmic relationship between measurements of the least shaft circumference of the femur/ tibiotarsus and body masses. Predictions of body mass, as well as the analysis of anatomical evidence, are used to infer that *Macranhinga paranensis* would have probably had a powered flapping flight and an aquatic locomotion similar to that of cormorants. The morphology of the pelvis and the hind-limb would have allowed *Macranhinga paranensis* to catch fishes by means of pursuit-diving rather than stalking them in an anhinga-like manner. As determined by adaptations mainly of tarsometatarsal morphology, the species had well developed perching and climbing abilities.

Key words: Aves, Anhingidae, *Macranhinga*, Paleobiology, Miocene, Argentina.

Jorge Ignacio Noriega [cidnoriega@infoshopde.com.ar], Centro de Investigaciones Científicas y de Transferencia Tecnológica - CONICET. Dr. Materi y España, (3105) Diamante, provincia de Entre Ríos, Argentina.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.