

Unique anatomy of lagomorph calcaneus

Ann R. Bleefeld and W. J. Bock *Acta Palaeontologica Polonica* 47 (1), 2002: 181-183

The mammalian order Lagomorpha (hares, rabbits, pikas) comprises two families (Ochotonidae and Leporidae; McKenna and Bell 1998; Nowak 1999), and Recent members have a nearly world-wide distribution (Hoffman 1993; Nowak 1999). Detailed examination of the pedal morphology of extant and fossil lagomorphs revealed a unique channel (the "calcaneal canal") running diagonally through the lagomorph calcaneus. The ancientness, ubiquity, and appearance of the calcaneal canal in all, including the earliest recognized lagomorph calcanea, and its absence from the pedes of other mammalian taxa, may indicate a long evolutionary separation of lagomorphs from other, previously suggested lagomorph relatives.

Ann R. Bleefeld [annrebek@amnh.org], Division of Paleontology, Ameri—can Museum of Natural History, 79th Street@Central ParkWest New York, New York 10024 USA; Walter J. Bock [wb4@columbia.edu], Department of Biological Sciences, Columbia University, 1200 Amsterdam Avenue, Mail Box 5521, New York, 10027 USA.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see <u>creativecommons.org</u>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

