

## A new heterosoricid shrew from the lowermost Oligocene of Europe

Richard Smith and Lars W. van den Hoek Ostende

*Acta Palaeontologica Polonica* 51 (2), 2006: 381-384

The earliest Oligocene faunas of Europe are characterised by a large number of Asian immigrants. One of the classical sites that shows the faunal change after this so-called “Grande Coupure” is Hoogbutsel in Belgium (MP 21). Recently a new locality from the lowermost Oligocene was discovered in Belgium, Boutersem TGV. Here, we describe a new heterosoricid (Mammalia, Soricomorpha), *Belgicasorex ramboeri* gen. et sp. nov., from these lowermost Oligocene localities. We assume that, like the erinaceid *Tetracus nanus* Aymard, 1846 and the nyctitheriid *Oligonyctia hoffmani* Smith, 2004, *Belgicasorex ramboeri* was one of the Asian immigrants that entered Europe after the “Grande Coupure”.

Richard Smith [[richardsmithpal@hotmail.com](mailto:richardsmithpal@hotmail.com)], Département de Paléontologie, Institut royal des Sciences naturelles de Belgique, rue Vautier 29, B-1000 Bruxelles, Belgium; Lars W. van den Hoek Ostende [[Hoek@Naturalis.nl](mailto:Hoek@Naturalis.nl)], Nationaal Natuurhistorisch Museum, Naturalis, Postbus 9517, NL-2300 RA Leiden, The Netherlands.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



[Full text \(163.7 kB\)](#)