

Antiquity of the substrate choice among acmaeid limpets from Late Cretaceous chemosynthesis-based communities

Robert G. Jenkins, Andrzej Kaim, and Yoshinori Hikida
Acta Palaeontologica Polonica 52 (2), 2007: 369-373

Two Campanian methane seep sites in the Nakagawa area of Hokkaido (northern Japan) yield fossils of the limpet genera *Serradonta* and *Bathyacmaea* that appear to have had the same substrate preference as do their modern counterparts. *Serradonta* cf. *vestimentifericola* was a species having an elongated and strongly compressed shell adapted to living on vestimentiferan tubes, like its modern relatives. *Bathyacmaea* cf. *subnipponica* was an acmaeid with a relatively elongated shell but with a more rounded aperture than *Serradonta* and thus apparently attached to small hard objects other than worm tubes. One *Bathyacmaea* specimen was found attached *in situ* to an ataphrid gastropod shell. The restricted present-day distribution of *Serradonta* possibly reflects its spreading route exclusively through the hot vent and cold seep communities settled by vestimentiferans.

Key words: Gastropoda, paleoecology, chemosynthetic community, methane-seep, Cretaceous, Hokkaido, Japan.

Robert Jenkins robert@um.u-tokyo.ac.jp, University Museum and Department of Earth and Planetary Science, University of Tokyo, Tokyo 113-0033, Japan ; Andrzej Kaim kaim@twarda.pan.pl, Institute of Paleobiology, Polish Academy of Sciences, PL-00-818 Warszawa, Poland and Department of Earth and Planetary Science, University of Tokyo, Tokyo 113-0033, Japan ; Yoshinori Hikida nmhikida@coral.ocn.ne.jp, Nakagawa Museum of Natural History, Hokkaido 068-0835, Japan.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(421.4 kB\)](#)