

Microconchid tubeworms from the Jurassic of England and France

Olev Vinn and Paul D. Taylor

Acta Palaeontologica Polonica 52 (2), 2007: 391-399

The Bajocian tubeworm *Spirorbis midfordensis*, previously regarded as a spirorbid polychaete, is reinterpreted as a microconchid and assigned to *Punctaconchus* gen. nov. along with two new species, *Punctaconchus ampliporus* sp. nov. (Toarcian?, Aalenian-Bathonian), the type species of the new genus, and *Punctaconchus palmeri* sp. nov. (Bathonian). Microconchids are a mostly Palaeozoic group of tubeworms that are probably more closely related to modern lophophorate phyla than they are to polychaetes. *Punctaconchus*, the youngest unequivocal microconchid, is characterised by having large pores (punctae) penetrating the tube wall, which has a fibrous or platy lamellar microstructure, and ripplemark-like transverse ridges on the tube interior. In both morphology and ecology it is a remarkable homeomorph of the polychaete *Spirorbis*.

Key words: Microconchida, morphology, Jurassic, England, France.

Olev Vinn olev.vinn@ut.ee, Geoloogia Instituut, University of Tartu, Vanemuise 46, 51014 Tartu, Estonia ; Paul D. Taylor p.taylor@nhm.ac.uk, Department of Palaeontology, Natural History Museum, Cromwell Road, London SW7 5BD, U.K.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.