

The hind limb skeleton and cursorial adaptations of the Plio-Pleistocene rabbit *Hypolagus beremendensis*

Łucja Fostowicz-Frelik

Acta Palaeontologica Polonica 52 (3), 2007: 447-476

Hypolagus beremendensis, a representative of the Archaeolaginae, was one of the most abundant and widespread leporids in the Plio-Pleistocene of Europe. The vast accumulations of skeletal remains from the Polish Pliocene sites (Węże 1, Rębiełice Królewskie 1 and 2, and Kadzielnia 1) yielded thousands of bones representing almost all skeletal regions. The detailed hind limb morphology of *Hypolagus beremendensis* is presented in comparison with five extant leporids (*Lepus europaeus*, *Oryctolagus cuniculus*, *Pentalagus furnessi*, *Sylvilagus floridanus*, and *S. brasiliensis*), which represent a wide range of locomotor adaptations. The UPGMA analysis of 98 metric characters places *Hypolagus beremendensis* next to the leporine rabbits. *Hypolagus beremendensis* has the os coxae, femur, and talus most similar to *P. furnessi*, tibia and calcaneus to the leporine rabbits, and the structure of foot to *Lepus*. The elongation of the foot and tibiofibular segment in relation to the femur indicates an advanced cursorial adaptation and a relatively steep jump. The similarities in the proximal segments (os coxae and femur) between *Hypolagus* and *Pentalagus* highlight the conservative morphology of this region in the Leporidae.

Key words: Lagomorpha, Leporidae, *Hypolagus beremendensis*, hind limb, functional morphology, cursorial adaptations, Neogene.

Łucja Fostowicz-Frelik lfost@twarda.pan.pl, Institute of Paleobiology Polish Academy of Sciences, ul. Twarda 51/55, PL-00-818 Warszawa, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(1,320.0 kB\)](#)