

## **Boine snake *Bavarioboa* from the Oligocene/Miocene of eastern Turkey with comments on connections between European and Asiatic snake faunas**

Zbigniew Szyndlar and Izzet Hoşgör

*Acta Palaeontologica Polonica* 57 (3), 2012: 667-671 doi: <http://dx.doi.org/10.4202/app.2011.0075>

Fossil remains of the extinct boine snake *Bavarioboa*, thus far known exclusively from several localities of western and central Europe, are reported for the first time outside Europe. The new fossil record is from the Mendikdere Formation in easternmost Turkey, dated Late Oligocene to Early Miocene. The finding provides strong evidence of links connecting ophidian faunas of Europe and southwestern Asia in the past, and confirms the supposition that Anatolia may have had close terrestrial connections with Europe around the Oligocene/Miocene boundary.

Zbigniew Szyndlar [[szyndlar@isez.pan.krakow.pl](mailto:szyndlar@isez.pan.krakow.pl)], Institute of Systematics and Evolution of Animals of the Polish Academy of Sciences, Sławkowska 17, 31-016 Kraków, Poland; Izzet Hoşgör [[Izzet.Hosgor@viking-intl.com](mailto:Izzet.Hosgor@viking-intl.com)], Viking International, Akmerkez, Block B Floor 5-6, Nispetiye Caddesi, 34330 Etiler-İstanbul, Turkey.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.