

## The fulturae ("compound synapticulae"), their structure and reconsideration of their systematic value


Gabriel A. Gill

*Acta Palaeontologica Polonica* 25 (3-4), 1980: 301-310

The study of the structure of fulturae in Recent *Cycloseria cyclolites* and various species of *Fungia* reveals an upward growth of fibers gradually diverging from the centre outwards. The distal end of each fultura is rounded and rhythmic secretion leaves a succession of horseshoe figures on the outer surface. In their structure, the fulturae differ basically from "simple" synapticulae and from the design which had been ascribed to them in literature. The twofold notion of synapticulae ("simple" and "compound"), until now considered as fundamental distinctive character for the suborder Fungilda, should be abandoned. The suborder Fungilda, as delineated in recent classifications, seems to include at least two apparent, non-related groups: one common in Tertiary and Recent forms which bear fulturae and show a septal structure similar to the *Montlivaltia* type (e.g. *Cycloseris*) and the other including pennular corals (mainly Mesozoic) with or without "simple" synapticulae.

**Key words:** Hexacoralla, Fungildae, skeletal microstructure.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(2,257.5 kB\)](#)