

A basal archosauriform from the early Triassic of Poland

Magdalena Borsuk-Białynicka and Susan E. Evans

Acta Palaeontologica Polonica 48 (4), 2003: 649-652

Basal Archosauriformes had a wide geographic distribution through the Lower to Middle Triassic. *Osmolskina czatkowiensis* gen. et sp. nov. from Early Olenekian karst deposits at Czatkowice, west of Cracow, provides the first record from Poland. The reconstructed skull and attributed postcranial elements show a morphology closely resembling that of the Early Anisian African genus *Euparkeria* Broom, 1913, while differing at generic level. Both genera display the same mosaic of plesiomorphic and apomorphic character states, but share no unique apomorphic character state. They might thus be combined in the family Euparkeriidae Huene, 1920, but could also constitute two plesions of the same grade lying just below the Archosauria + Proterochampsidae node. Currently, Euparkeriidae remains monotypic because no other genus can be assigned to it with confidence. Until this problem is resolved, the term “euparkeriid” essentially denotes a grade of Lower to Middle Triassic non-archosaurian archosauriforms that are more derived than proterosuchid grade taxa, but lack the specializations of either erythrosuchids or proterochampsids. They were probably Pangaeian in their distribution.

Magdalena Borsuk-Białynicka [borsuk.b@twarda.pan.pl], Instytut Paleobiologii Polskiej Akademii Nauk, ul. Twarda 51/55, PL-00-818 Warszawa, Poland; Susan E. Evans [ucgasue@ucl.ac.uk], Department of Anatomy and Developmental Biology, University College, Gower Street, London, WC1E 6BT, England.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.