

Revision of Middle Ordovician orthoceratacean nautiloids from Baltoscandia

Björn Kröger

Acta Palaeontologica Polonica 49 (1), 2004: 57-74

The fauna of the Baltoscandic Orthocerataceae is important because it fills a documentary gap between the Lower Ordovician fauna of North America and the upper Middle Ordovician fauna of China and North America. A revision of the Orthoceratidae, Geisonoceratidae, and Arionoceratidae is given on a material of more than 450 specimens. Intraspecific and ontogenetic variations were observed. The following new taxa are erected: *Plagiostomoceras fragile* sp. nov., *Archigeisonoceras repplingense* sp. nov., *Archigeisonoceras picus* sp. nov., *Archigeisonoceras folkeslundense* sp. nov., *Nilssonoceras latisiphonatum* gen. et sp. nov., *Kinnekulloceras kinnekulense* gen. et sp. nov., and *Arionoceras lotskirkense* sp. nov. The genus *Archigeisonoceras* is described for the first time in the Baltoscandic area and the oldest occurrence of the genus *Arionoceras* is documented from the Middle Ordovician of Baltoscandia. It is shown that the endosiphuncular deposits of the Middle Ordovician orthocerataceans are highly variable and widespread in the apical parts of the phragmocone.

Key words: Cephalopoda, Nautiloidea, Orthoceratacea, Ordovician, Baltoscandia.

Björn Kröger [buxcreau@gmx.de], Geologisch-Paläontologisches Institut und Museum, Universität Hamburg, Bundesstr. 55, D-20146 Hamburg, Germany.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

