

## The dentition of a eugeneodontiform shark from the Lower Pennsylvanian of Derbyshire, UK

Michał Ginter

*Acta Palaeontologica Polonica* 63 (4), 2018: 725-735 doi:<https://doi.org/10.4202/app.00533.2018>

The anterior part of a medium-sized shark surrounded by hundreds of ostracod shells was found at the end of the last century in a dark limestone nodule from the Kinderscoutian (Bashkirian, Pennsylvanian) near Carsington, Derbyshire (England, UK). The shark is a caseodontoid eugeneodontiform, most probably belonging to *Campodus agassizianus*. Its dentition is of the crushing type, highly heterodont, but the symphyseal tooth whorl, typical of the most of eugeneodontiforms, was not found. The teeth are symmetrical labio-lingually which is unique in this group. The analysis of the available data leads to the conclusion that neither the tooth whorls of “*Campodus variabilis*” sensu Eastman, nor the mandibular dentition of “*Agassizodus variabilis*” sensu St. John and Worthen represent the genus *Campodus* and that these specimens deserve a new, probably common, name.

**Key words:** Chondrichthyes, Eugeneodontiformes, *Campodus*, teeth, Carboniferous, Bashkirian, UK, England.

Michał Ginter [[m.ginter@uw.edu.pl](mailto:m.ginter@uw.edu.pl)], Faculty of Geology, University of Warsaw, Al. Żwirki i Wigury 93, 02-089 Warszawa, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

