

## A new capybara from the late Miocene of San Juan Province, Argentina, and its phylogenetic implications

Esperanza Cerdeño, María E. Pérez, Cecilia M. Deschamps, and Víctor H. Contreras

*Acta Palaeontologica Polonica* 64 (1), 2019: 199-212 doi:<https://doi.org/10.4202/app.00544.2018>

A new hydrochoerine rodent, *Cardiatherium calingastaense* sp. nov. (Caviidae), is described based on the specimen INGEO-PV 87. It was recovered from the late Miocene Las Flores Formation, cropping out at the Puchuzum locality, San Juan Province, Argentina. The new species is based on a particular combination of characters, among which the following can be highlighted: p4 with internal fissures equally deep as in *Cardiatherium paranense*, secondary external fissure as in *Cardiatherium patagonicum*, and lacking the fifth internal fissure and supernumerary internal fissure, as in *C. paranense*; m3 with a conspicuous labial column in the posterior ramus of the second prism; very deep primary and secondary external fissures in upper cheek teeth, the former producing a labial strong step-shaped profile in M2; sagittal crest on the parietals; bullae small in ventral view; scars of the origin of the masseter medialis muscle with an anterior projection up to the level of the incisive foramen and the maxilla-premaxilla suture. The phylogenetic analysis supports the taxonomic proposal of creating a new species of *Cardiatherium* and shows *C. calingastaense* sp. nov. as the sister group of the other species of the genus. The lineage leading to the clade *Cardiatherium* + largest capybaras would have originated at least during the Chasicoan SALMA (early late Miocene). *Cardiatherium calingastaense* sp. nov. adds to the previous record of *Cardiatherium chasicoense* and *Cardiatherium paranense* in the late Miocene of San Juan and Mendoza provinces, respectively, thus increasing the diversity of capybaras in central-west Argentina.

**Key words:** Mammalia, Rodentia, Caviidae, *Cardiatherium*, systematics, phylogeny, late Miocene, Argentina.

Esperanza Cerdeño [[espe@mendoza-conicet.gob.ar](mailto:espe@mendoza-conicet.gob.ar)], Instituto Argentino de Nivología, Glaciología y Ciencias Ambi-entales (IANIGLA), Centro Científico Tecnológico, CONICET Mendoza, Av. Ruiz Leal s/n, 5500, Mendoza, Argentina. María E. Pérez [[mperez@mef.org.ar](mailto:mperez@mef.org.ar)], Museo Paleontológico Egidio Feruglio, Av. Fontana 140, U9100GYO, Trelew, Argentina. Cecilia M. Deschamps [[ceci@fcnym.unlp.edu.ar](mailto:ceci@fcnym.unlp.edu.ar)], Comisión de Investigaciones Científicas de la provincia de Buenos Aires, Argentina; División

Paleontología Vertebrados, Unidades de Investigación Anexo Museo de La Plata, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, 122 y 60, 1900, La Plata, Argentina. Víctor H. Contreras [[vcontre@unsj-cuim.edu.ar](mailto:vcontre@unsj-cuim.edu.ar)], Instituto de Geología Dr. Emiliano P. Aparicio (INGEO) y Departamento Geología, FCEF-N, Universidad Nacional de San Juan, Ignacio de la Roza y Meglioli, 5400 Rivadavia, San Juan, Argentina.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(634.1 kB\)](#) |

 [Supplementary file \(231.9 kB\)](#)