

## Considerations on Middle and Upper Devonian *Thamnophyllidae* soshkina in Poland

Maria Rózkowska

*Acta Palaeontologica Polonica* 02 (2-3), 1957: 81-153

Corals with horse-shoe dissepiments and resting on them trabecular fans, formerly included by the writer in the subfamily *Pachyphyllinae* Stumm, are now referred to the family *Thamnophyllidae* Soshkina. The systematic division of this family is given. An analysis of correlation between the number of septa and the diameter of calice or of tabularium (n/d or n/t) in genera *Thamnophyllum* Penecke, *Macgeea* Webster, *Pexiphyllum* Walther and *Pachyphyllum* Edward & Haime has made possible the determination of the range of individual variability and the elucidation of evolutionary tendencies within the quoted genera. The coefficient of correlation for particular species is of diagnostic value and may constitute their stratigraphical index. It has been established that species previously described and referred (Rózkowska, 1953, 1956) to the genera - *Pseudoacervularia* Schluter and *Pachyphyllum* Edwards & Haime should be assigned to genus *Pachyphyllum*. Within this genus four groups of species, with peculiar, similar coefficients of correlation, should be differentiated. Conclusions concerning speciation have been deduced from biometrical treatment and from ontogenic and morphological studies. A description is given of the phyletic and chronological evolution of *Thamnophyllidae*. In addition to material from the Holy Cross Mts. specimens collected at Dębnik and in the Sudeten were used by the writer for the present work.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org/)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

