

## Miocene coniferous woods of the Polish Carpathian Foredeep

Marzena Klusek

*Acta Palaeontologica Polonica* 59 (3), 2014: 697-708 doi: <http://dx.doi.org/10.4202/app.2011.0158>


This paper presents the results of analysis of the Miocene woods collected from three sampling sites in the Polish part of the Carpathian Foredeep. The location in the stratigraphic profile and the state of fossil preservation indicate that the woods were transported by waters flowing from highland or from mountainous regions and then deposited and fossilised in delta areas. Fragments of 16 lignified or silicified woods were collected for anatomical research. The fossil woods were classified as *Piceoxylon piceae*, *Juniperoxylon pachyderma*, *Cupressinoxylon canadense* and *Cupressinoxylon polonicum*. Disturbance zones observed in *Cupressinoxylon polonicum* are interpreted as frost rings. Frost rings suggest that despite the documented existence of warm and humid climatic conditions during the Miocene period, the temperatures in mountainous areas could temporarily drop below freezing point.

**Key words:** Coniferophyta, fossil woods, palaeoecology, wood anatomy, Miocene, Poland.

Marzena Klusek [[marzena.klusek@boku.ac.at](mailto:marzena.klusek@boku.ac.at)], BOKU University of Natural Resources and Applied Life Sciences Vienna, University Research Center Tulln, Konrad Lorenz Strasse 24, 3430 Tulln an der Donau, Austria.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(1,174.4 kB\)](#) |

 [Supplementary file \(190.0 kB\)](#)



