

## New teeth of nodosaurid ankylosaur from the Lower Cretaceous of Southern England

William T. Blows and Kerri Honeysett

*Acta Palaeontologica Polonica* 59 (4), 2014: 835-841 doi: <http://dx.doi.org/10.4202/app.2012.0131>

We present new nodosaurid teeth from the Valanginian of Bexhill, Sussex and the Barremian of the Isle of Wight, the first from the Lower Cretaceous of the United Kingdom. Teeth found during the mid-1800s from the Valanginian and ascribed to the nodosaurid *Hylaeosaurus* are probably from sauropod dinosaurs. The Isle of Wight tooth could possibly be referred to *Polacanthus foxii*, the teeth of which are unknown. These new English nodosaurid teeth are similar to those of North American and European Jurassic to Late Cretaceous nodosaurids, especially the American *Gastonia*, *Texasetes*, *Mymoorapelta*, *Gargoyleosaurus*, and the European *Hungarosaurus*.

**Key words:** Dinosauria, Nodosauridae, *Polacanthus*, ankylosaur, Valanginian, Barremian, Cretaceous, England.

William T. Blows [[w.t.blows@city.ac.uk](mailto:w.t.blows@city.ac.uk)] Department of Applied Biological Sciences, City University, London, EC1A 7QN, England;  
Kerri Honeysett [[kerrihoneysett@googlemail.com](mailto:kerrihoneysett@googlemail.com)], 10 Camperdown Street, Bexhill on Sea, East Sussex, TN39 5BE, England.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

