

Ultrastructure of the jaws of the fossil and Recent Eunicida (Polychaeta)

Grażyna Mierzejewska and Piotr Mierzejewski

Acta Palaeontologica Polonica 23 (3), 1978: 317-339

The ultrastructure of the jaws of Ordovician, Silurian, Permian and Recent Eunicida (Polychaeta) have been studied by transmission and scanning electron microscopy. Primary and secondary zones have been distinguished in the ultrastructure of the jaws. The primary zone is formed of the pharyngeal cuticle containing an orthogonal network of collagen fibrils, while the secondary zone, underlying the primary one, is developed of inner part of the pharyngeal cuticle devoid of a regular network of collagen fibrils. The two zones are pierced by microvilli. The jaws of placognath Eunicida are structured of the primary one only. The Eunicida with highly organized jaw apparatus have a strongly developed secondary zone underlying the primary one. A preliminary correlation of evolutionary changes in the anatomy and ultrastructure of the jaws of Eunicida has been conducted. It has also been shown that the ultrastructural studies of the scolecodonts were not very useful for taxonomic purposes.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(2,215.2 kB\)](#)