

Rugose corals from the Devonian Kowala Formation of the Holy Cross Mountains

Tomasz Wrzosek

Acta Palaeontologica Polonica 37 (2-4), 1992: 217-254

Rugose corals of the Givetian to Lower Frasnian Kowala limestone Formation in the environs of Chęciny, SW Holy Cross Mts and in its age equivalents in the Silesian-Cracow region of Southern Poland represent five distinct assemblages of restricted time-and-space distribution. Within the *Stringocephalus* Beds the high diversity *Pseudohexagonaria(?) laxa* assemblage indicates open-shelf conditions whereas low diversity *Temnophyllum occidentale* assemblage represents restricted conditions. The transgressive Jaźwica Mbr. locally contains diversified and cosmopolitan *Acanthophyllum* sp. n. fauna. Following temporally coral assemblages, i.e., *Disphyllum* (lower Sitkówka and Chęciny Beds) and *Macgeea-Thamnophyllum* (Kadzielnia Mbr, upper Sitkówka Beds) are mostly biostromes of branching corals of low taxonomical diversity typical for restricted relatively setting, rather unfavorable for rugosans. Exceptional are two *Hexagonaria* horizons with common massive colonies. *Diffusolasma* gen. nov., *Sociophyllum severiacum* sp. nov., *Temnophyllum zamkowsae* sp. nov. and *Hexagonaria hexagona kowalae* subsp. nov. are proposed as the new taxa.

Key words: Rugosa, Devonian, Holy Cross Mts, Kowala Formation, taxonomy, stratigraphical distribution.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

