

Maastrichtian *Ceratisepia* and Mesozoic cuttlebone homeomorphs

Roger A. Hewitt and John W.M. Jagt

Acta Palaeontologica Polonica 44 (3), 1999: 305-326

The phylogenetics of potential Mesozoic ancestors of cuttlefish of a restricted order Sepiida von Zittel, 1895 (superorder Decabrachia Boettger, 1952) is reviewed. Microstructural studies of Mesozoic homeomorphs of cuttlebones (*Pearceiteuthis* gen. n., *Loligosepia*, *Trachyteuthis*, *Actinosepia*) are consistent with their assignment to the superorder Octobrachia Fioroni, 1981. The discovery of an embryonic *Ceratisepia* shell in the upper Maastrichtian of the Netherlands, indicates that true Sepiida did have a pre-Cenozoic origin. Cretaceous decabrachs of the order Spirulida Stolley, 1919 do not show evidence of the dorso-anterior shell growth vectors seen in Cenozoic spirulids, sepiids and octibrachs. Separate origins of the Sepiida and Spirulida within Cretaceous diplobelinid belemnites is still the most attractive hypothesis. *Ceratisepia vanknippenbergi* sp. n. from the upper Maastrichtian of the Netherlands and *Pearceiteuthis buyi* gen. et sp. n. from the Callovian of England are described.

Key words: Cephalopods, cuttlefish, ontogeny, biomineralization, Mesozoic.

Roger A. Hewitt, 12 Fairfield Road, Eastwood, Leigh-on-Sea, Essex SS9 5SB,
United Kingdom. John W.M. Jagt [mail@nhmmaastricht.nl], Natuurhistorisch Museum Maastricht, P.O.
Box 882, NL-6200 AW Maastricht, the Netherlands.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.