

A new discosauriscid seymouriamorph tetrapod from the Lower Permian of Moravia, Czech Republic

Jozef Klembara

Acta Palaeontologica Polonica 50 (1), 2005: 25-48

A new genus and species, *Makowskia laticephala* gen. et sp. nov., of seymouriamorph tetrapod from the Lower Permian deposits of the Boskovice Furrow in Moravia (Czech Republic) is described in detail, and its cranial reconstruction is presented. It is placed in the family Discosauriscidae (together with *Discosauriscus* and *Ariekanerpeton*) on the following character states: short preorbital region; rounded to oval orbits positioned mainly in anterior half of skull; otic notch dorsoventrally broad and anteroposteriorly deep; rounded to oval ventral scales. *Makowskia* is distinguished from other Discosauriscidae by the following characters: nasal bones equally long as broad; interorbital region broad; prefrontal-postfrontal contact lies in level of frontal mid-length (only from *D. pulcherrimus*); maxilla deepest at its mid-length; suborbital ramus of jugal short and dorsoventrally broad with long anterodorsal-posteroventral directed lacrimal-jugal suture; postorbital anteroposteriorly short and lacks elongated posterior process; ventral surface of basioccipital smooth; rows of small denticles placed on distinct ridges and intervening furrows radiate from place immediately laterally to articular portion on ventral surface of palatal ramus of pterygoid (only from *D. pulcherrimus*); oblique anterior margin of transverse flange of pterygoid directed anteromedially-posterolaterally; cultriform process of parasphenoid relatively short and slightly rounded; ventral surface of the posterior plate of parasphenoid heavily sculptured (only from *D. pulcherrimus* and *Ariekanerpeton*); distal ends of fourth and fifth presacral ribs distinctly anteroposteriorly broadened, and extend into the hook-like, posteriorly directed processes; shaft of ?last caudal rib anteroposteriorly broadened; posterior stem of interclavicle narrows anteriorly and posteriorly from broadened mid-length section.

Key words: Skeletal anatomy, Discosauriscidae, Seymouriamorpha, Permian, Boskovice Furrow, Moravia, Czech Republic.

Jozef Klembara [klembara@fns.uniba.sk], Comenius University in Bratislava, Faculty of Natural Sciences, Mlynská dolina, 842 15 Bratislava, Slovak Republic.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(878.8 kB\)](#)