

## Ostracods and facies of the Lower and Middle Frasnian at Devils Gate in Nevada: Relationship to the Alamo Event.

Jean-Georges Casier, Ivan Berra, Ewa Olempska, Charles Sandberg, and Alain Pr  at  
*Acta Palaeontologica Polonica* 51 (4), 2006: 813-828

In order to document the Alamo Event and to investigate its influence on shallow-marine environments, we undertook a study of ostracods, conodonts, and analysis of the sedimentology of the lower member of the type Devils Gate Limestone. Six major carbonate microfacies (MF1-MF6) ranging from open-marine environments below storm wave base to pre-evaporitic supratidal lagoons were recognized. The sedimentological study detected no important sedimentological changes during the Alamo Event; only an influx of detrital material and lithoclasts indicate that an unusual event had occurred. Ostracods are generally rare or absent in the lower member of the Devils Gate Limestone, and only 2,000 carapaces, valves and fragments were extracted; from these some 26 taxa were identified. Two new species, *Voronina* ? *eureka* and *Serenida dorsoplicata* are proposed. The ostracods belong to the Eifelian Mega-Assemblage and their distribution was influenced by strong salinity variations. Because of the rarity and low diversity of ostracods and conodonts in samples collected from the lower part of the lower member of the Devils Gate Limestone it is not adequate to demonstrate conclusively an extinction event close to the Alamo Event Bed. Nevertheless the greater abundance and diversity of ostracods above this bed seems to indicate that the Alamo Event did not result in significant extinction of ostracod taxa in this shallow water setting. The ostracod fauna present in the lower member of the Devils Gate Limestone suggests faunal exchanges between Nevada and the Russian Platform via the Western Canadian platform.

**Key words:** Ostracoda, conodonts, sedimentology, palaeoecology, Frasnian, Alamo Event, Pilot Basin, Nevada.

Jean-Georges Casier [casier@naturalsciences.be](mailto:casier@naturalsciences.be), D  partement de Pal  ontologie, Institut royal des Sciences naturelles de Belgique, 29, rue Vautier, B   1000, Bruxelles, Belgique; Ewa Olempska [olempska@twarda.pan.pl](mailto:olempska@twarda.pan.pl), Instytut Paleobiologii PAN, ul. Twarda 51/55, PL   00   818 Warszawa, Poland; Ivan Berra [iberra@ulb.ac.be](mailto:iberra@ulb.ac.be) and Alain Pr  at [apreat@ulb.ac.be](mailto:apreat@ulb.ac.be), D  partement des Sciences de la Terre et de l'Environnement, Universit   libre de Bruxelles CP 160, 50, Av. F.D. Roosevelt, B   1050, Bruxelles, Belgique; Charles Sandberg, U.S. [sandberg@usgs.gov](mailto:sandberg@usgs.gov), Geological Survey, box 25046, Mail Stop 939, Federal Center, Denver, Colorado 80225, USA.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(899.6 kB\)](#)